

Datasheet: MCA1578

Description:	MOUSE ANTI HUMAN CD34
Specificity:	CD34 CLASS III
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	581
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting		▪		

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Cynomolgus monkey</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid.
Preparation	Purified IgG prepared by affinity chromatography on Protein A.
Buffer Solution	Phosphate buffered saline

Preservative <0.1% Sodium Azide (NaN₃)
Stabilisers 0.2% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.1 mg/ml

External Database Links

UniProt:
[P28906](#) [Related reagents](#)

Entrez Gene:
[947](#) CD34 [Related reagents](#)

RRID AB_1125259

Specificity

Mouse anti Human CD34 antibody, clone 581 recognizes the human CD34 cell surface antigen also known as Hematopoietic progenitor cell antigen CD34. Clone 581 binds to an epitope of human CD34 resistant to neuraminidase, chymopapain and glycoprotease treatments. The binding epitopes are classified according to their resistance to enzymatic degradation and antibodies assigned according to this and cross inhibition studies. Clone 581 is classified as class III ([Lanza *et al.* 1999](#)).

CD34 is a 385 amino acid pro-peptide with the 31 amino acid signal peptide cleaved to yield the mature form of CD34. CD34 is a single pass type I ~105-120 kDa highly glycosylated transmembrane glycoprotein, expressed by hematopoietic stem cells and small vessel endothelial cells. CD34 contains a number of sialylated residues and has a mucin-like structure ([Simmons *et al.* 1992](#)). A truncated form of the CD34 molecule has also been identified lacking the carboxy-terminal region of the intracellular sequence, also bearing a sequence switch in the remaining cytoplasmic region ([Nakamura *et al.* 1993](#)). Mouse anti Human CD34 antibody, clone 581 is expected to react with both isoforms of the CD34 molecule as it recognizes an epitope located in the extracellular region of human CD34.

Mouse anti Human CD34 antibody, clone 581 has been used successfully for the immunohistochemical detection of stem-like cells present in hepatoblastoma lesions on cryostat sections ([Fiegel *et al.* 2004](#)).

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

Histology Positive Control Tissue Tonsil, Bone marrow

References

1. Civin, C.I. *et al.* (1984) Antigenic analysis of hematopoiesis. III. A hematopoietic progenitor cell surface antigen defined by a monoclonal antibody raised against KG-1a cells. [J Immunol. 133 \(1\): 157-65.](#)
2. Fiegel, H.C. *et al.* (2004) Stem-like cells in human hepatoblastoma. [J Histochem Cytochem. 52 \(11\): 1495-501.](#)
3. Maumus, M. *et al.* (2011) Native human adipose stromal cells: localization, morphology and phenotype. [Int J Obes \(Lond\). 35 \(9\): 1141-53.](#)

4. Patel, J. *et al.* (2013) Prospective surface marker-based isolation and expansion of fetal endothelial colony-forming cells from human term placenta. [Stem Cells Transl Med. 2 \(11\): 839-47.](#)
5. Park, J.Y. *et al.* (2020) rhBMP-2 Pre-Treated Human Periodontal Ligament Stem Cell Sheets Regenerate a Mineralized Layer Mimicking Dental Cementum. [Int J Mol Sci. 21 \(11\): 3767.](#)
6. Kim, S.H. *et al.* (2019) Forkhead box O1 (FOXO1) controls the migratory response of Toll-like receptor (TLR3)-stimulated human mesenchymal stromal cells. [J Biol Chem. 294 \(21\): 8424-37.](#)

Further Reading 1. Yoshino, N. *et al.* (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of Cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. [Exp. Anim. 49: 97-110.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

Worldwide Tel: +44 (0)1865 852 700

Europe Tel: +49 (0) 89 8090 95 21

To
find a

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M389307:210806'

Printed on 29 Aug 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)