

Datasheet: MCA1580FT

BATCH NUMBER 020614

Description:	HAMSTER ANTI MOUSE CD69:FITC
Specificity:	CD69
Other names:	AIM
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	H1.2F3
Isotype:	IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat - 1/10
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugat	ed to Fluorescein Isoth	niocyanate Isomer 1
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide	(NaN ₃)	
Approx. Protein Concentrations	IgG concentration 0.5	5 mg/ml.	

Immunogen External Database	Murine dendritic epidermal Y245 cell line.
Links	UniProt:
	P37217 Related reagents
	Entrez Gene:
	12515 Cd69 Related reagents
RRID	AB_2075078
Specificity	Hamster anti Mouse CD69 antibody, clone H1.2F3 recognizes the murine CD69 cell surface antigen, an ~85 kDa dimeric glycoprotein. CD69 is expressed rapidly on the surface of T cells, B cells and NK cells following activation.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
	The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (<u>BUF041A/B</u>).
References	 Yokoyama, W.M. <i>et al.</i> (1988) Characterization of a cell surface-expressed disulfide-linked dimer involved in murine T cell activation. <u>J Immunol. 141 (2): 369-76.</u> Hickman, H.D. <i>et al.</i> (2008) Direct priming of antiviral CD8+ T cells in the peripheral interfollicular region of lymph nodes. <u>Nat Immunol. 9 (2): 155-65.</u> Chalifour, A. <i>et al.</i> (2004) Direct bacterial protein PAMP recognition by human NK cells involves TLRs and triggers alpha-defensin production. <u>Blood. 104 (6): 1778-83.</u> Cohen-sfady, M. <i>et al.</i> (2005) Heat shock protein 60 activates B cells via the TLR4-MyD88 pathway. <u>J Immunol. 175 (6): 3594-602.</u> Iwashiro, M. <i>et al.</i> (2001) Immunosuppression by CD4+ regulatory T cells induced by chronic retroviral infection. <u>Proc Natl Acad Sci U S A. 98 (16): 9226-30.</u>

cells over time. <u>J Immunol. 170 (4): 2022-9.</u>

7. Kvakan, H. *et al.* (2009) Regulatory T cells ameliorate angiotensin II-induced cardiac damage. <u>Circulation. 119 (22): 2904-12.</u>

8. Marshall-clarke, S. *et al.* (2003) A differential requirement for phosphoinositide 3-kinase reveals two pathways for inducible upregulation of major histocompatibility complex class II molecules and CD86 expression by murine B lymphocytes. <u>Immunology</u>. 109 (1): 102-8.

6. Karrer, U. et al. (2003) Memory inflation: continuous accumulation of antiviral CD8+ T

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

 Guarantee
 12 months from date of despatch

 Health And Safety Information
 Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1580FT

 Regulatory
 For research purposes only

Related Products

Recommended Negative Controls

HAMSTER (ARMENIAN) IgG NEGATIVE CONTROL:FITC (MCA2356F)

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A)
MOUSE SEROBLOCK FcR (BUF041B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739

Email: antibody_sales_us@bio-rad.com

Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365502:200529'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint