

## Datasheet: MCA1580FT

**BATCH NUMBER 162960**

<b>Description:</b>	HAMSTER ANTI MOUSE CD69:FITC
<b>Specificity:</b>	CD69
<b>Other names:</b>	AIM
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	H1.2F3
<b>Isotype:</b>	IgG
<b>Quantity:</b>	0.1 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse						
<b>Product Form</b>	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
<b>Max Ex/Em</b>	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					
<b>Buffer Solution</b>	Phosphate buffered saline						
<b>Preservative Stabilisers</b>	<0.1% Sodium Azide (NaN <sub>3</sub> )						
<b>Approx. Protein Concentrations</b>	IgG concentration 0.5 mg/ml.						

**Immunogen** Murine dendritic epidermal Y245 cell line.

---

**External Database**

**Links**

**UniProt:**

[P37217](#)    [Related reagents](#)

**Entrez Gene:**

[12515](#) Cd69    [Related reagents](#)

---

**RRID**

AB\_2075078

---

**Specificity**

**Hamster anti Mouse CD69 antibody, clone H1.2F3** recognizes the murine CD69 cell surface antigen, an ~85 kDa dimeric glycoprotein. CD69 is expressed rapidly on the surface of T cells, B cells and NK cells following activation.

---

**Flow Cytometry**

Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#)).

---

**References**

1. Yokoyama, W.M. *et al.* (1988) Characterization of a cell surface-expressed disulfide-linked dimer involved in murine T cell activation. [J Immunol. 141 \(2\): 369-76.](#)
  2. Hickman, H.D. *et al.* (2008) Direct priming of antiviral CD8+ T cells in the peripheral interfollicular region of lymph nodes. [Nat Immunol. 9 \(2\): 155-65.](#)
  3. Chalifour, A. *et al.* (2004) Direct bacterial protein PAMP recognition by human NK cells involves TLRs and triggers alpha-defensin production. [Blood. 104 \(6\): 1778-83.](#)
  4. Cohen-sfady, M. *et al.* (2005) Heat shock protein 60 activates B cells via the TLR4-MyD88 pathway. [J Immunol. 175 \(6\): 3594-602.](#)
  5. Iwashiro, M. *et al.* (2001) Immunosuppression by CD4+ regulatory T cells induced by chronic retroviral infection. [Proc Natl Acad Sci U S A. 98 \(16\): 9226-30.](#)
  6. Karrer, U. *et al.* (2003) Memory inflation: continuous accumulation of antiviral CD8+ T cells over time. [J Immunol. 170 \(4\): 2022-9.](#)
  7. Kvakan, H. *et al.* (2009) Regulatory T cells ameliorate angiotensin II-induced cardiac damage. [Circulation. 119 \(22\): 2904-12.](#)
  8. Marshall-clarke, S. *et al.* (2003) A differential requirement for phosphoinositide 3-kinase reveals two pathways for inducible upregulation of major histocompatibility complex class II molecules and CD86 expression by murine B lymphocytes. [Immunology. 109 \(1\): 102-8.](#)
- 

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA1580FT>  
10040

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL:FITC \(MCA2356F\)](#)

### Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M390140:210825'

**Printed on 18 Jan 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)