

Datasheet: MCA1580FT

Description:	HAMSTER ANTI MOUSE CD69:FITC
Specificity:	CD69
Other names:	AIM
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	H1.2F3
Isotype:	IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)						
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml.						
Immunogen	Murine dendritic epidermal Y245 cell line.						

External Database Links	UniProt: P37217 Related reagents Entrez Gene: 12515 Cd69 Related reagents
RRID	AB_2075078
Specificity	Hamster anti Mouse CD69 antibody, clone H1.2F3 recognizes the murine CD69 cell surface antigen, an ~85 kDa dimeric glycoprotein. CD69 is expressed rapidly on the surface of T cells, B cells and NK cells following activation.
Flow Cytometry	<p>Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.</p> <p>The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).</p>
References	<ol style="list-style-type: none"> 1. Yokoyama, W.M. <i>et al.</i> (1988) Characterization of a cell surface-expressed disulfide-linked dimer involved in murine T cell activation. J Immunol. 141 (2): 369-76. 2. Hickman, H.D. <i>et al.</i> (2008) Direct priming of antiviral CD8+ T cells in the peripheral interfollicular region of lymph nodes. Nat Immunol. 9 (2): 155-65. 3. Chalifour, A. <i>et al.</i> (2004) Direct bacterial protein PAMP recognition by human NK cells involves TLRs and triggers alpha-defensin production. Blood. 104 (6): 1778-83. 4. Cohen-sfady, M. <i>et al.</i> (2005) Heat shock protein 60 activates B cells via the TLR4-MyD88 pathway. J Immunol. 175 (6): 3594-602. 5. Iwashiro, M. <i>et al.</i> (2001) Immunosuppression by CD4+ regulatory T cells induced by chronic retroviral infection. Proc Natl Acad Sci U S A. 98 (16): 9226-30. 6. Karrer, U. <i>et al.</i> (2003) Memory inflation: continuous accumulation of antiviral CD8+ T cells over time. J Immunol. 170 (4): 2022-9. 7. Kvakon, H. <i>et al.</i> (2009) Regulatory T cells ameliorate angiotensin II-induced cardiac damage. Circulation. 119 (22): 2904-12. 8. Marshall-clarke, S. <i>et al.</i> (2003) A differential requirement for phosphoinositide 3-kinase reveals two pathways for inducible upregulation of major histocompatibility complex class II molecules and CD86 expression by murine B lymphocytes. Immunology. 109 (1): 102-8.
Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf

Related Products

Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL:FITC \(MCA2356F\)](#)

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M390140:210825'

Printed on 06 Jan 2022
