

## Datasheet: MCA1582T

<b>Description:</b>	MOUSE ANTI HUMAN CD83
<b>Specificity:</b>	CD83
<b>Other names:</b>	HB15
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	HB15e
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	20 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			1/500 - 1/1000
Immunohistology - Paraffin (1)	▪			1/50
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1) This product requires protein digestion pre-treatment of paraffin sections e.g. trypsin or pronase.**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	<p>Reacts with: Cynomolgus monkey, Chimpanzee, Baboon, Rhesus Monkey, Tasmanian Devil, Sheep</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
<b>Product Form</b>	Purified IgG - liquid

<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% sodium azide (NaN <sub>3</sub> )
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Cos cells transfected with HB15 cDNA.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">Q01151</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">9308</a>    CD83    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_324789
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Human CD83 antibody, clone HB15e</b> recognizes the human CD83 cell surface antigen, a 40-45 kDa glycoprotein expressed by peripheral blood dendritic cells. Peripheral lymphocytes can be induced to express very low levels of CD83 after culture in agents such as Con A or PHA.</p> <p>In immunohistology CD83 is shown to be expressed strongly by interfollicular interdigitating reticulum cells and more weakly by cells within germinal centres. CD83 is also expressed by Langerhan's cells in the skin. The CD83 antigen is a 186-amino-acid single-chain glycoprotein. This molecule is a member of the immunoglobulin superfamily and is composed of an extracellular V-type Ig-like single domain, a transmembrane region, and a short, 40-amino-acid cytoplasmic tail. CD83 antigen undergoes extensive post-translational glycosylation, since the determined Mr is twice the predicted size of the core protein (<a href="#">Zhou et al. 1992</a>).</p> <p>However, CD83+ cells have a unique cell surface immuno-phenotype that does not correlate with that of T cells, B cells, NK cells, or cells of the myelomonocytic lineage (<a href="#">Zhou et al. 1995</a>). CD83+ cells co-express the highest levels of MHC class II molecules, when compared with other leucocyte lineages. They also co-express T cell markers (CD2, CD5), B cell markers (CD40, CD78), myeloid cell markers (CD13, CD33, CD36), cytokine receptors as well as other cell surface molecules (<a href="#">Zhou et al. 1995</a>) and <a href="#">Zhou and Tedder 1995</a>).</p>

**Flow Cytometry** Use 10µl of the suggested working dilution to label 10<sup>6</sup>cells in 100µl

**Histology Positive Control Tissue** Human tonsil

#### References

1. Zhou, L.J. *et al.* (1992) A novel cell-surface molecule expressed by human interdigitating reticulum cells, Langerhans cells, and activated lymphocytes is a new member of the Ig superfamily. [J Immunol. 149 \(2\): 735-42.](#)
2. Zhou, L.J. & Tedder, T.F. (1995) Human blood dendritic cells selectively express CD83, a member of the immunoglobulin superfamily. [J Immunol. 154 \(8\): 3821-35.](#)
3. Zhou, L.J. & Tedder, T.F. (1995) A distinct pattern of cytokine gene expression by human CD83+ blood dendritic cells. [Blood. 86 \(9\): 3295-301.](#)
4. Denniston, A.K. *et al.* (2011) Endogenous Cortisol and TGF- $\beta$  in Human Aqueous Humor Contribute to Ocular Immune Privilege by Regulating Dendritic Cell Function. [J Immunol. 186:305-11.](#)
5. Schlossman, S.F., *et al.* Eds. Engel, P. *et al.* (1995) 'CD83 Workshop report' in Leucocyte Typing V, White Cell Differentiation Antigens, Oxford University Press pp. 693-5.
6. Yoshino, N. *et al.* (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. [Exp Anim. 49 \(2\): 97-110.](#)
7. Hesselink, D.A. *et al.* (2005) The effects of renal transplantation on circulating dendritic cells. [Clin Exp Immunol. 140: 384-93.](#)
8. Hovden, A.O. *et al.* (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. [BMC Immunol. ; 12:2.](#)
9. Ifergan, I. *et al.* (2008) The blood-brain barrier induces differentiation of migrating monocytes into Th17-polarizing dendritic cells. [Brain. 131: 785-99.](#)
10. Walker, J.G. *et al.* (2007) Characterisation of a dendritic cell subset in synovial tissue which strongly expresses Jak/STAT transcription factors from patients with rheumatoid arthritis. [Ann Rheum Dis. 66: 992-9.](#)
11. Köller, M. *et al.* (2004) Phenotypic and functional deficiencies of monocyte-derived dendritic cells in systemic lupus erythematosus (SLE) patients. [Int Immunol. 16: 1595-604.](#)
12. Denniston, A.K. *et al.* (2012) Aqueous humor suppression of dendritic cell function helps maintain immune regulation in the eye during human uveitis. [Invest Ophthalmol Vis Sci. 53 \(2\): 888-96.](#)
13. Shikotra, A. *et al.* (2012) Increased expression of immunoreactive thymic stromal lymphopoietin in patients with severe asthma. [J Allergy Clin Immunol. 129: 104-11.e1-9.](#)
14. Sprater, F. *et al.* (2012) Expression of ESE-3 Isoforms in Immunogenic and Tolerogenic Human Monocyte-Derived Dendritic Cells [PLoS One. 7: e49577.](#)
15. Silk, K.M. *et al.* (2012) Rapamycin conditioning of dendritic cells differentiated from human ES cells promotes a tolerogenic phenotype. [J Biomed Biotechnol. 2012: 172420.](#)
16. Howson, L.J. *et al.* (2014) Identification of dendritic cells, B cell and T cell subsets in Tasmanian devil lymphoid tissue; evidence for poor immune cell infiltration into devil facial tumors. [Anat Rec \(Hoboken\). 297: 925-38.](#)
17. Eren, U. *et al.* (2016) The several elements of intestinal innate immune system at the beginning of the life of broiler chicks. [Microsc Res Tech. 79 \(7\): 604-14.](#)
18. Wang, P. *et al.* (2016) Distribution and expression profiles of dendritic cell subpopulations in human bladder cancer. [Int J Clin Exp Pathol 9\(7\):7180-7.](#)

19. Van Vré, E.A. *et al.* (2011) Immunohistochemical characterisation of dendritic cells in human atherosclerotic lesions: possible pitfalls. [Pathology. 43 \(3\): 239-47.](#)
20. Duan, Y.G. *et al.* (2016) Characterisation of dendritic cell subsets in chronically inflamed human epididymis. [Andrologia. 48 \(4\): 431-40.](#)
21. Arya, S. *et al.* (2019) Quantitative proteomic changes in LPS-activated monocyte-derived dendritic cells: A SWATH-MS study. [Sci Rep. 9 \(1\): 4343.](#)
22. Pérez-caballero, R. *et al.* (2018) Comparative dynamics of peritoneal cell immunophenotypes in sheep during the early and late stages of the infection with *Fasciola hepatica* by flow cytometric analysis. [Parasit Vectors. 11 \(1\): 640.](#)
23. Yildiz, M. *et al.* (2019) Histological and immunohistochemical studies of the proximal caecum and caecal tonsils of quail (*Coturnix coturnix japonica*). [Anat Histol Embryol. 48 \(5\): 476-85.](#)
24. Eren, U. *et al.* (2022) TLR2 and TLR4 molecules and antigen-presenting cell compositions in cecal tonsils of broiler chicks (*Gallus gallus domesticus*.) in the first two weeks of the post-hatch period. [Anat Histol Embryol. 51 \(1\): 125-35.](#)
25. Barbieri, S. *et al.* (2022) Does smoking habit affect dendritic cell expression in oral squamous cell carcinoma? [Braz Oral Res. 36: e044.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

**Regulatory** For research purposes only

## Related Products

### Recommended Secondary Antibodies

- |   |   |
|---|---|
| Goat Anti Mouse IgG (STAR77...)         | <a href="#">HRP</a>   |
| Rabbit Anti Mouse IgG (STAR12...)       | <a href="#">RPE</a>   |
| Goat Anti Mouse IgG (STAR70...)         | <a href="#">FITC</a>  |
| Goat Anti Mouse IgG IgA IgM (STAR87...) | <a href="#">Alk. Phos.</a> , <a href="#">HRP</a>  |
| Rabbit Anti Mouse IgG (STAR9...)        | <a href="#">FITC</a>  |
| Goat Anti Mouse IgG (STAR76...)         | <a href="#">RPE</a>   |
| Goat Anti Mouse IgG (H/L) (STAR117...)  | <a href="#">Alk. Phos.</a> , <a href="#">DyLight@488</a> , <a href="#">DyLight@550</a> ,<br><a href="#">DyLight@650</a> , <a href="#">DyLight@680</a> , <a href="#">DyLight@800</a> ,<br><a href="#">FITC</a> , <a href="#">HRP</a> |
| Rabbit Anti Mouse IgG (STAR13...)       | <a href="#">HRP</a>   |
| Goat Anti Mouse IgG (Fc) (STAR120...)   | <a href="#">FITC</a> , <a href="#">HRP</a>  |

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

**North & South** Tel: +1 800 265 7376

**Worldwide**

Tel: +44 (0)1865 852 700

**Europe**

Tel: +49 (0) 89 8090 95 21

**America** Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M414884:221214'

**Printed on 15 Dec 2022**

---

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)