

## Datasheet: MCA1584

**BATCH NUMBER 153749**

<b>Description:</b>	MOUSE ANTI HUMAN CD158b
<b>Specificity:</b>	CD158b
<b>Other names:</b>	KIR2DL3
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	GL183
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.2 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - liquid
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	NK cell clone E57 ( <a href="#">Moretta et al. 1985</a> ).

<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P43628</a> <a href="#">Related reagents</a>
	<b>Entrez Gene:</b> <a href="#">3804</a> KIR2DL3 <a href="#">Related reagents</a>
<b>Synonyms</b>	CD158B2, KIRCL23, NKAT2
<b>RRID</b>	AB_2265256
<b>Fusion Partners</b>	Spleen cells from immunized Balb/c mice were fused with cells of the mouse P3UI myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Human CD158b antibody, clone GL183</b> recognizes human Killer cell immunoglobulin-like receptor 2DL3, also known as CD158b, KIR-023GB, MHC class I NK cell receptor, p58 natural killer cell receptor clone CL-6 or Natural killer-associated transcript 2. CD158b is a 341 amino acid, ~58 kDa single pass type-1 transmembrane glycoprotein containing two <a href="#">Ig-like C2-type</a> domains. expressed by a subset of NK cells.</p> <p>This antibody also recognizes a ~50 kDa molecule in some NK clones, which is highly homologous to p58.2 in the extracellular domain, but has a shorter cytoplasmic tail (<a href="#">Moretta et al. 1985</a>). Both molecules are members of the newly described natural killer cell receptor family.</p> <p>CD158b functions as a receptor specific for HLA Class I molecules, including <a href="#">Cw3</a> and related HLA-C alleles. Mouse anti Human CD158b antibody, clone GL183 can restore the lysis by human NK clones of otherwise lysis protected targets expressing Cw3.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Moretta, A. <i>et al.</i> (1990) A novel surface antigen expressed by a subset of human CD3-CD16+ natural killer cells. Role in cell activation and regulation of cytolytic function. <a href="#">J Exp Med. 171 (3): 695-714.</a></li> <li>2. Moretta, A. <i>et al.</i> (1993) P58 molecules as putative receptors for major histocompatibility complex (MHC) class I molecules in human natural killer (NK) cells. Anti-p58 antibodies reconstitute lysis of MHC class I-protected cells in NK clones displaying different specificities. <a href="#">J Exp Med. 178 (2): 597-604.</a></li> <li>3. Moretta, A. <i>et al.</i> (1995) Existence of both inhibitory (p58) and activatory (p50) receptors for HLA-C molecules in human natural killer cells. <a href="#">J Exp Med. 182 (3): 875-84.</a></li> <li>4. Pridgeon, C. <i>et al.</i> (2003) Natural killer cells in the synovial fluid of rheumatoid arthritis patients exhibit a CD56bright,CD94bright,CD158negative phenotype. <a href="#">Rheumatology (Oxford). 42 (7): 870-8.</a></li> <li>5. Marget, M. <i>et al.</i> (2005) A HLA-Cw6 specific single-chain antibody fragment (scFv) recognizing a natural killer cell receptor epitope <a href="#">Mol Immunol. 42: 643-9.</a></li> <li>6. Poggi, A. <i>et al.</i> (2005) Regulation of gammadelta T cell survival by soluble HLA-I: involvement of CD8 and activating killer Ig-like receptors. <a href="#">Eur J Immunol. 35: 2670-8.</a></li> <li>7. Borhis, G. <i>et al.</i> (2013) A peptide antagonist disrupts NK cell inhibitory synapse</li> </ol>

- formation. [J Immunol. 190 \(6\): 2924-30.](#)
8. Poggi, A. *et al.* (2005) Patients with paroxysmal nocturnal hemoglobinuria have a high frequency of peripheral-blood T cells expressing activating isoforms of inhibiting superfamily receptors. [Blood. 106: 2399-408.](#)
9. Valés-Gómez, M. *et al.* (2003) Expression of the UL16 glycoprotein of Human Cytomegalovirus protects the virus-infected cell from attack by natural killer cells. [BMC Immunol. 4:4.](#)
10. Spaggiari, G.M. *et al.* (2002) Soluble HLA class I molecules induce natural killer cell apoptosis through the engagement of CD8: evidence for a negative regulation exerted by members of the inhibitory receptor superfamily. [Blood. 99: 1706-14.](#)
11. Warren, H.S. *et al.* (2001) Biphasic response of NK cells expressing both activating and inhibitory killer Ig-like receptors. [Int Immunol. 13: 1043-52.](#)
12. Ghio, M. *et al.* (2009) Soluble HLA-I-mediated secretion of TGF-beta1 by human NK cells and consequent down-regulation of anti-tumor cytolytic activity. [Eur J Immunol. 39: 3459-68.](#)
13. Spaggiari, G.M. *et al.* (2003) IFN-gamma production in human NK cells through the engagement of CD8 by soluble or surface HLA class I molecules. [Eur J Immunol. 33: 3049-59.](#)
14. Bachelet, I. *et al.* (2005) The inhibitory receptor IRp60 (CD300a) is expressed and functional on human mast cells. [J Immunol. 175: 7989-95.](#)
15. Zimmer, J. *et al.* (1998) Activity and phenotype of natural killer cells in peptide transporter (TAP)-deficient patients (type I bare lymphocyte syndrome). [J Exp Med. 187: 117-22.](#)
16. Castriconi, R. *et al.* (2009) NK cells recognize and kill human glioblastoma cells with stem cell-like properties. [J Immunol. 182 \(6\): 3530-9.](#)
17. Ghio, M. *et al.* (2009) Soluble HLA-I-mediated secretion of TGF-beta1 by human NK cells and consequent down-regulation of anti-tumor cytolytic activity. [Eur J Immunol. 39 \(12\): 3459-68.](#)
18. Naiyer, M.M. *et al.* (2017) KIR2DS2 recognizes conserved peptides derived from viral helicases in the context of HLA-C. [Sci Immunol. 2 \(15\) \[Epub ahead of print\].](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1584>  
10040

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M365512:200529'

Printed on 01 May 2024