

Datasheet: MCA1590PET

BATCH NUMBER INN1608

Description:	MOUSE ANTI HUMAN CD40:RPE
Specificity:	CD40
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	LOB7/6
Isotype:	IgG2a
Quantity:	25 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Species Cross Reactivity	Reacts with: Dog N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute in 0.25 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Buffer Solution	Phosphate buffered saline		

Preservative	0.09% Sodium Azide
Stabilisers	1% Bovine Serum Albumin
Immunogen	CD40 Ig(Fc) fusion protein containing the EC region of human CD40 and Fc region of human IgG.
External Database Links	<p>UniProt: P25942 Related reagents</p> <p>Entrez Gene: 958 CD40 Related reagents</p>
Synonyms	TNFRSF5
RRID	AB_2275836
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
Specificity	<p>Mouse anti Human CD40 antibody, clone LOB7/6 recognizes the human CD40 cell surface antigen, a 48kDa glycoprotein expressed by B lymphocytes and weakly by some monocytes.</p> <p>CD40 is involved in the process of B cell selection in germinal centres and is vital in T cell-B cell interactions.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Quadbeck, B. <i>et al.</i> (2002) Maturation of thyroidal dendritic cells in Graves' disease. Scand J Immunol. 55 (6): 612-20. 2. Kirsch, B. M. <i>et al.</i> (2005) The active metabolite of leflunomide, A77 1726, interferes with dendritic cell function. Arthritis Res. Ther. 7: R694-R703. 3. Cheadle, E. <i>et al.</i> (2003) <i>Mycobacterium bovis</i> bacillus Calmette-Guerin-infected dendritic cells potently activate autologous T cells via a B7 and interleukin-12-dependent mechanism. Immunology.108: 79-88. 4. Carpenter, E.L. <i>et al.</i> (2009) Activation of human B cells by the agonist CD40 antibody CP-870,893 and augmentation with simultaneous toll-like receptor 9 stimulation. J Transl Med. 7: 93. 5. Garcia-Nieto, S. <i>et al.</i> (2010) Laminin and Fibronectin Treatment Leads to Generation of Dendritic Cells with Superior Endocytic Capacity. PLoS ONE. 5: 1-10. 6. Wang, Y.S. <i>et al.</i> (2007) Characterization of canine monocyte-derived dendritic cells with phenotypic and functional differentiation. Can J Vet Res. 71: 165-74. 7. Vlachoyiannopoulos, P.G. <i>et al.</i> (2004) Anti-CD40 antibodies in antiphospholipid syndrome and systemic lupus erythematosus. Thromb Haemost. 92: 1303-11. 8. Leigh, J.E. <i>et al.</i> (2006) Characterization of the immune status of CD8+ T cells in oral lesions of human immunodeficiency virus-infected persons with oropharyngeal Candidiasis. Clin Vaccine Immunol. 13: 678-83.

9. Newman, K.C. *et al.* (2006) Cross-talk with myeloid accessory cells regulates human natural killer cell interferon-gamma responses to malaria. [PLoS Pathog. 2: e118.](#)
10. Kuijf, M.L. *et al.* (2010) TLR4-mediated sensing of *Campylobacter jejuni* by dendritic cells is determined by sialylation. [J Immunol. 185: 748-55.](#)
11. Huizinga R *et al.* (2015) Innate Immunity to *Campylobacter jejuni* in Guillain-Barré Syndrome. [Ann Neurol. 78 \(3\): 343-54.](#)
12. Yildirim C *et al.* (2015) Galectin-2 induces a proinflammatory, anti-arteriogenic phenotype in monocytes and macrophages. [PLoS One. 10 \(4\): e0124347.](#)
13. Brencicova, E. *et al.* (2017) Interleukin-10 and prostaglandin E2 have complementary but distinct suppressive effects on Toll-like receptor-mediated dendritic cell activation in ovarian carcinoma. [PLoS One. 12 \(4\): e0175712.](#)
14. Silk, K.M. *et al.* (2012) Rapamycin conditioning of dendritic cells differentiated from human ES cells promotes a tolerogenic phenotype. [J Biomed Biotechnol. 2012: 172420.](#)
15. Tischer, S. *et al.* (2011) Heat shock protein 70/peptide complexes: potent mediators for the generation of antiviral T cells particularly with regard to low precursor frequencies. [J Transl Med. 9: 175.](#)
16. Unosson, J. *et al.* (2021) Acute cardiovascular effects of controlled exposure to dilute Petrodiesel and biodiesel exhaust in healthy volunteers: a crossover study. [Part Fibre Toxicol. 18 \(1\): 22.](#)
17. Milhau, N. *et al.* (2020) *In vitro.* evaluations on canine monocyte-derived dendritic cells of a nanoparticles delivery system for vaccine antigen against *Echinococcus granulosus.* [PLoS One. 15 \(2\): e0229121.](#)
18. Uetz-von Allmen, E *et al.* (2021) CAL-1 as Cellular Model System to Study CCR7-Guided Human Dendritic Cell Migration. [Front Immunol. 12: 702453.](#)

Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #20487 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1590PET20487>

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:RPE \(MCA929PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M375344:210104'

Printed on 21 Nov 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)