

Datasheet: MCA1614A488

Description:	MOUSE ANTI HUMAN CD55:Alexa Fluor® 488
Specificity:	CD55
Other names:	DAF
Format:	ALEXA FLUOR® 488
Product Type:	Monoclonal Antibody
Clone:	67
Isotype:	IgG1
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Alexa Fluor® 488 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®488	495	519
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml		
Immunogen	K562 cells		
External Database Links	UniProt:		
	P08174	Related reagents	

Entrez Gene:[1604](#) CD55 [Related reagents](#)**Synonyms**

CR, DAF

Specificity

Mouse anti Human CD55 antibody, clone 67 recognizes the human CD55 cell surface antigen, a GPI linked molecule also known as decay accelerating factor (DAF). CD55 is expressed by a wide range of cell types.

CD55 is the complement regulatory protein, decay accelerating factor (DAF) ([Lublin and Atkinson 1989](#)). Human CD55 is a ~70 kDa glycoprotein (in erythrocytes) anchored in the membrane by glycosylphosphatidylinositol tail. In other cells the apparent molecular weight is somewhat larger. It has a substantial content of O-glycans, and also on N-glycan. DAF binds to activated C4b or C3b complement fragments on the cell surface, preventing the assembly and accelerating the decay of both classical and alternative pathways. DAF carries the Cromer related blood group antigens.

DAF has a wide distribution on cells in non-haematopoietic tissues, particularly epithelium and is found at the fetal-maternal interface in placenta ([Holmes et al. 1990](#) and [Yang et al. 2009](#)). Soluble forms of DAF are found, for example, in plasma, saliva and urine ([Medof et al. 1987](#)). The antigen on erythrocytes is pronase and chymotrypsin sensitive, but resistant to trypsin.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul. Please note: Bio-Rad do not recommend the use of this reagent to stain erythrocytes

References

1. Hadam, M.R. (1989) In Leucocyte Typing IV: White Cell Differentiation Antigens. Edited by Knapp, W. *et al.* Oxford University Press pp 694-697.
2. Holmes, C.H. *et al.* (1990) Preferential expression of the complement regulatory protein decay accelerating factor at the fetomaternal interface during human pregnancy. [J Immunol. 144 \(8\): 3099-105.](#)
3. O'Brien, D.P. *et al.* (2009) Regulation of the Helicobacter pylori cellular receptor decay-accelerating factor. [J Biol Chem. 283: 23922-30.](#)
4. Yang, P. *et al.* (2009) Expression and modulation of RPE cell membrane complement regulatory proteins. [Invest Ophthalmol Vis Sci. 50: 3473-81.](#)
5. van de Sande, M.G. *et al.* (2011) Different stages of rheumatoid arthritis: features of the synovium in the preclinical phase. [Ann Rheum Dis. 70: 772-7.](#)
6. Mo, B. *et al.* (2006) ECC-1 cells: a well-differentiated steroid-responsive endometrial cell line with characteristics of luminal epithelium. [Biol Reprod. 75: 387-94.](#)
7. Araten, D.J. *et al.* (2005) A quantitative measurement of the human somatic mutation rate. [Cancer Res. 65: 8111-7.](#)
8. de Launay, D. *et al.* (2010) Silencing the expression of Ras family GTPase homologues decreases inflammation and joint destruction in experimental arthritis. [Am J Pathol. 177: 3010-24.](#)
9. Gheorghie, K.R. *et al.* (2011) Prostaglandin E2 synthesizing enzymes in rheumatoid arthritis B cells and the effects of B cell depleting therapy on enzyme expression. [PLoS One. ;6: e16378.](#)
10. Kraan, M.C. *et al.* (2004) T cells, fibroblast-like synoviocytes, and granzyme B+ cytotoxic cells are associated with joint damage in patients with recent onset rheumatoid arthritis. [Ann Rheum Dis. 63: 483-8.](#)
11. van Holten, J. *et al.* (2005) A multicentre, randomised, double blind, placebo controlled phase II study of subcutaneous interferon beta-1a in the treatment of patients with active rheumatoid arthritis. [Ann Rheum Dis. 64 \(1\): 64-9.](#)
12. Abreu, J.R. *et al.* (2009) The Ras guanine nucleotide exchange factor RasGRF1 promotes matrix metalloproteinase-3 production in rheumatoid arthritis synovial tissue. [Arthritis Res Ther. 11\(4\):R121.](#)

13. Thurlings, R.M. *et al.* (2008) Synovial tissue response to rituximab: mechanism of action and identification of biomarkers of response. [Ann Rheum Dis. 67 \(7\): 917-25.](#)
14. Vos, K. *et al.* (2007) Early effects of rituximab on the synovial cell infiltrate in patients with rheumatoid arthritis. [Arthritis Rheum. 56 \(3\): 772-8.](#)
15. Edginton S *et al.* (2016) Effects of Rituximab and Infliximab Treatment on Carboxypeptidase B and Its Substrates in RA Synovium. [J Rheumatol. 43 \(5\): 846-54.](#)

Further Reading	<p>1. Lublin, D.M. & Atkinson, J.P. (1989) Decay-accelerating factor: biochemistry, molecular biology, and function. Annu Rev Immunol. 7: 35-58.</p> <p>2. Daniels, G. (1989) Cromer-related antigens--blood group determinants on decay-accelerating factor. Vox Sang. 56 (4): 205-11.</p>
------------------------	---

Storage	<p>Store at +4°C or at -20°C if preferred. This product should be stored undiluted.</p> <p>Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
----------------	--

Guarantee	12 months from date of despatch
------------------	---------------------------------

Acknowledgements	<p>This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchased product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com</p>
-------------------------	--

Health And Safety Information	<p>Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf</p>
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 488 \(MCA928A488\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	<p>Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com</p>	Worldwide	<p>Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com</p>	Europe	<p>Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com</p>
----------------------------------	--	------------------	--	---------------	--

'M365532:200529'

