

## Datasheet: MCA1614GA

<b>Description:</b>	MOUSE ANTI HUMAN CD55
<b>Specificity:</b>	CD55
<b>Other names:</b>	DAF
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	67
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/25
Immunohistology - Frozen	▪			1/100 - 1/1000
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	K562 cells

**External Database  
Links**

**UniProt:**

[P08174](#) [Related reagents](#)

**Entrez Gene:**

[1604](#) CD55 [Related reagents](#)

---

**Synonyms**

CR, DAF

---

**RRID**

AB\_324645

---

**Specificity**

**Mouse anti Human CD55 antibody, clone 67** recognizes the human CD55 cell surface antigen, a GPI linked molecule also known as decay accelerating factor (DAF). CD55 is expressed by a wide range of cell types.

CD55 is the complement regulatory protein, decay accelerating factor (DAF) ([Lublin and Atkinson 1989](#)). Human CD55 is a ~70 kDa glycoprotein (in erythrocytes) anchored in the membrane by glycosylphosphatidylinositol tail. In other cells the apparent molecular weight is somewhat larger. It has a substantial content of O-glycans, and also on N-glycan. DAF binds to activated C4b or C3b complement fragments on the cell surface, preventing the assembly and accelerating the decay of both classical and alternative pathways. DAF carries the Cromer related blood group antigens.

DAF has a wide distribution on cells in non-haematopoietic tissues, particularly epithelium and is found at the fetal-maternal interface in placenta ([Holmes et al. 1990](#) and [Yang et al. 2009](#)). Soluble forms of DAF are found, for example, in plasma, saliva and urine ([Medof et al. 1987](#)). The antigen on erythrocytes is pronase and chymotrypsin sensitive, but resistant to trypsin.

---

**Flow Cytometry**

Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul. Please note: Bio-Rad do not recommend the use of this reagent to stain erythrocytes

---

**Histology Positive  
Control Tissue**

Human Tonsil

---

**References**

1. Hadam, M.R. (1989) In Leucocyte Typing IV: White Cell Differentiation Antigens. Edited by Knapp, W. *et al.* Oxford University Press pp 694-697.
2. Holmes, C.H. *et al.* (1990) Preferential expression of the complement regulatory protein decay accelerating factor at the fetomaternal interface during human pregnancy. [J Immunol. 144 \(8\): 3099-105.](#)
3. O'Brien, D.P. *et al.* (2009) Regulation of the Helicobacter pylori cellular receptor decay-accelerating factor. [J Biol Chem. 283: 23922-30.](#)
4. Yang, P. *et al.* (2009) Expression and modulation of RPE cell membrane complement regulatory proteins. [Invest Ophthalmol Vis Sci. 50: 3473-81.](#)
5. van de Sande, M.G. *et al.* (2011) Different stages of rheumatoid arthritis: features of the synovium in the preclinical phase. [Ann Rheum Dis. 70: 772-7.](#)
6. Mo, B. *et al.* (2006) ECC-1 cells: a well-differentiated steroid-responsive endometrial cell line with characteristics of luminal epithelium. [Biol Reprod. 75: 387-94.](#)
7. Araten, D.J. *et al.* (2005) A quantitative measurement of the human somatic mutation rate. [Cancer Res. 65: 8111-7.](#)
8. de Launay, D. *et al.* (2010) Silencing the expression of Ras family GTPase homologues decreases inflammation and joint destruction in experimental arthritis. [Am J Pathol. 177: 3010-24.](#)
9. Gheorghe, K.R. *et al.* (2011) Prostaglandin E2 synthesizing enzymes in rheumatoid arthritis B cells and the effects of B cell depleting therapy on enzyme expression. [PLoS One. ;6: e16378.](#)
10. Kraan, M.C. *et al.* (2004) T cells, fibroblast-like synoviocytes, and granzyme B+ cytotoxic cells

are associated with joint damage in patients with recent onset rheumatoid arthritis. [Ann Rheum Dis. 63: 483-8.](#)

11. van Holten, J. *et al.* (2005) A multicentre, randomised, double blind, placebo controlled phase II study of subcutaneous interferon beta-1a in the treatment of patients with active rheumatoid arthritis. [Ann Rheum Dis. 64 \(1\): 64-9.](#)

12. Abreu, J.R. *et al.* (2009) The Ras guanine nucleotide exchange factor RasGRF1 promotes matrix metalloproteinase-3 production in rheumatoid arthritis synovial tissue. [Arthritis Res Ther.11\(4\):R121.](#)

13. Thurlings, R.M. *et al.* (2008) Synovial tissue response to rituximab: mechanism of action and identification of biomarkers of response. [Ann Rheum Dis. 67 \(7\): 917-25.](#)

14. Vos, K. *et al.* (2007) Early effects of rituximab on the synovial cell infiltrate in patients with rheumatoid arthritis. [Arthritis Rheum. 56 \(3\): 772-8.](#)

15. Edginton S *et al.* (2016) Effects of Rituximab and Infliximab Treatment on Carboxypeptidase B and Its Substrates in RA Synovium. [J Rheumatol. 43 \(5\): 846-54.](#)

---

<b>Further Reading</b>	1. Lublin, D.M. & Atkinson, J.P. (1989) Decay-accelerating factor: biochemistry, molecular biology, and function. <a href="#">Annu Rev Immunol. 7: 35-58.</a> 2. Daniels, G. (1989) Cromer-related antigens--blood group determinants on decay-accelerating factor. <a href="#">Vox Sang. 56 (4): 205-11.</a>
------------------------	--

---

<b>Storage</b>	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
----------------	---

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: 10040: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a>
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR8...)	<a href="#">DyLight®800</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

**North & South America** Tel: +1 800 265 7376  
Fax: +1 919 878 3751  
Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide** Tel: +44 (0)1865 852 700  
Fax: +44 (0)1865 852 739  
Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)  
'M365536:200529'

**Europe** Tel: +49 (0) 89 8090 95 21  
Fax: +49 (0) 89 8090 95 50  
Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

**Printed on 11 Aug 2020**

---

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)