

## Datasheet: MCA1615GA

<b>Description:</b>	MOUSE ANTI HUMAN CD54
<b>Specificity:</b>	CD54
<b>Other names:</b>	ICAM-1
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	15.2
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			1/50 - 1/100
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

#### Target Species

Human

#### Species Cross Reactivity

Reacts with: Pig

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

#### Product Form

Purified IgG - liquid

#### Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Human monocytes
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P05362</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">3383</a> ICAM1    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_324159
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse Sp2/0-Ag14 myeloma cell line
<b>Specificity</b>	<p><b>Mouse anti Human CD54 antibody, clone 15.2</b> recognizes the human CD54 cell surface antigen also known as intracellular Adhesion Molecule -1 (ICAM-1) or Major group rhinovirus receptor.</p> <p>CD54 is expressed by many cells following activation by inflammatory mediators. It is a 505 amino acid with an additional 27 amino acid signal peptide ~90 kDa single pass type I transmembrane glycoprotein bearing 5 Ig-like C2-type domains.</p> <p>Mouse anti Human CD54 antibody, clone 15.2 blocks CD54 function (<a href="#">Berendt <i>et al.</i> 1992</a>). Mouse anti Human CD54 antibody, clone 15.2 binds to an epitope on the N-terminal Ig-like domain within a region designated the L43 loop (<a href="#">Chakravorty and Craig 2005</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>Histology Positive Control Tissue</b>	Human Tonsil
<b>References</b>	<ol style="list-style-type: none"> <li>1. Dransfield, I. <i>et al.</i> (1992) Interaction of leukocyte integrins with ligand is necessary but not sufficient for function. <a href="#">J Cell Biol. 116:1527-35.</a></li> <li>2. Berendt, A. <i>et al.</i> (1992) The binding site on ICAM-1 for plasmodium falciparum-infected erythrocytes overlaps, but is distinct from, the LFA-1- binding site. <a href="#">Cell. 68: 71-81.</a></li> <li>3. Urquhart, P. <i>et al.</i> (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. <a href="#">J Pharmacol Exp Ther. 321 (2): 656-62.</a></li> <li>4. Baratin, M. <i>et al.</i> (2007) Dissection of the role of PfEMP1 and ICAM-1 in the sensing of <i>Plasmodium-falciparum</i>-infected erythrocytes by natural killer cells. <a href="#">PLoS One 2: e228.</a></li> </ol>

5. van Buul, J.D. *et al.* (2010) Inside-out regulation of ICAM-1 dynamics in TNF-alpha-activated endothelium. [PLoS One 5: e11336.](#)
6. Diaz-Romero, J. *et al.* (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect bona fide dedifferentiation rather than amplification of progenitor cells. [J Cell Physiol. 214: 75-83.](#)
7. Di Lorenzo, A. *et al.* (2011) Endothelial reticulon-4B (Nogo-B) regulates ICAM-1-mediated leukocyte transmigration and acute inflammation. [Blood. 117: 2284-95.](#)
8. Porter, J.C. and Hall, A. (2009) Epithelial ICAM-1 and ICAM-2 regulate the egression of human T cells across the bronchial epithelium. [FASEB J. 23: 492-502.](#)
9. Corvaisier, M. *et al.* (2005) V gamma 9V delta 2 T cell response to colon carcinoma cells. [J Immunol. 175: 5481-8.](#)
10. Horrocks, P. *et al.* (2005) PfEMP1 expression is reduced on the surface of knobless Plasmodium falciparum infected erythrocytes. [J Cell Sci. 118: 2507-18.](#)
11. Lozanoska-Ochser, B. *et al.* (2008) Expression of CD86 on human islet endothelial cells facilitates T cell adhesion and migration. [J Immunol. 181: 6109-16.](#)
12. Norling, L.V. *et al.* (2008) Inhibitory control of endothelial galectin-1 on *in vitro* and *in vivo* lymphocyte trafficking. [FASEB J. 22: 682-90.](#)
13. Baumer, Y. *et al.* (2011) Telomerase-based immortalization modifies the angiogenic/inflammatory responses of human coronary artery endothelial cells. [Exp Biol Med \(Maywood\). 236: 692-700.](#)
14. Lask, A. *et al.* (2011) TCR-independent killing of B cell malignancies by anti-third-party CTLs: the critical role of MHC-CD8 engagement. [J Immunol. 187 \(4\): 2006-14.](#)
15. Sommaggio, R. *et al.* (2012) Multiple Receptors Trigger Human NK Cell-Mediated Cytotoxicity against Porcine Chondrocytes. [J Immunol. 188: 2075-83.](#)
16. Murphy, A.J. *et al.* (2013) Anti-inflammatory functions of apolipoprotein a-I and high-density lipoprotein are preserved in trimeric apolipoprotein a-I. [J Pharmacol Exp Ther. 344: 41-9.](#)
17. Sumagin R *et al.* (2014) Transmigrated neutrophils in the intestinal lumen engage ICAM-1 to regulate the epithelial barrier and neutrophil recruitment. [Mucosal Immunol. 7 \(4\): 905-15.](#)
18. Sugden SM *et al.* (2017) HIV-1 Vpu Downmodulates ICAM-1 Expression, Resulting in Decreased Killing of Infected CD4<sup>+</sup> T Cells by NK Cells. [J Virol. 91 \(8\): pii: e02442-16.](#)
19. Lennartz, F. *et al.* (2015) Mapping the Binding Site of a Cross-Reactive *Plasmodium falciparum* PfEMP1 Monoclonal Antibody Inhibitory of ICAM-1 Binding. [J Immunol. 195 \(7\): 3273-83.](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR8...)	<a href="#">DyLight®800</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M365549:200529'

Printed on 18 Apr 2021