

Datasheet: MCA1615P647

Description:	MOUSE ANTI HUMAN CD54:RPE-Alexa Fluor® 647
Specificity:	CD54
Other names:	ICAM-1
Format:	RPE-ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	15.2
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Pig

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to RPE-Alexa Fluor 647 - lyophilized

Reconstitution

Reconstitute with 1.0 ml distilled water

Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
RPE-Alexa Fluor®647 488nm laser	496	667

RPE-Alexa Fluor®647
561nm laser

546

667

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
--------------------	---

Buffer Solution	Phosphate buffered saline
------------------------	---------------------------

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin 5% Sucrose
-------------------------------------	---

Immunogen	Human monocytes
------------------	-----------------

External Database Links	UniProt: P05362 Related reagents Entrez Gene: 3383 ICAM1 Related reagents
------------------------------------	--

Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse Sp2/0-Ag14 myeloma cell line
------------------------	---

Specificity	<p>Mouse anti Human CD54 antibody, clone 15.2 recognizes the human CD54 cell surface antigen also known as intracellular Adhesion Molecule -1 (ICAM-1) or Major group rhinovirus receptor.</p> <p>CD54 is expressed by many cells following activation by inflammatory mediators. It is a 505 amino acid with an additional 27 amino acid signal peptide ~90 kDa single pass type I transmembrane glycoprotein bearing 5 Ig-like C2-type domains.</p> <p>Mouse anti Human CD54 antibody, clone 15.2 blocks CD54 function (Berendt et al. 1992). Mouse anti Human CD54 antibody, clone 15.2 binds to an epitope on the N-terminal Ig-like domain within a region designated the L43 loop (Chakravorty and Craig 2005).</p>
--------------------	--

Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul
-----------------------	--

References	<ol style="list-style-type: none">1. Dransfield, I. <i>et al.</i> (1992) Interaction of leukocyte integrins with ligand is necessary but not sufficient for function. J Cell Biol. 116:1527-35.2. Berendt, A. <i>et al.</i> (1992) The binding site on ICAM-1 for plasmodium falciparum-infected erythrocytes overlaps, but is distinct from, the LFA-1- binding site. Cell. 68: 71-81.3. Urquhart, P. <i>et al.</i> (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. J Pharmacol Exp Ther. 321 (2): 656-62.4. Baratin, M. <i>et al.</i> (2007) Dissection of the role of PfEMP1 and ICAM-1 in the sensing of <i>Plasmodium-falciparum</i>-infected erythrocytes by natural killer cells. PLoS One 2: e228.5. van Buul, J.D. <i>et al.</i> (2010) Inside-out regulation of ICAM-1 dynamics in TNF-alpha-activated endothelium. PLoS One 5: e11336.6. Diaz-Romero, J. <i>et al.</i> (2008) Immunophenotypic changes of human articular
-------------------	--

- chondrocytes during monolayer culture reflect bona fide dedifferentiation rather than amplification of progenitor cells. [J Cell Physiol. 214: 75-83.](#)
7. Di Lorenzo, A. *et al.* (2011) Endothelial reticulon-4B (Nogo-B) regulates ICAM-1-mediated leukocyte transmigration and acute inflammation. [Blood. 117: 2284-95.](#)
 8. Porter, J.C. and Hall, A. (2009) Epithelial ICAM-1 and ICAM-2 regulate the egression of human T cells across the bronchial epithelium. [FASEB J. 23: 492-502.](#)
 9. Corvaisier, M. *et al.* (2005) V gamma 9V delta 2 T cell response to colon carcinoma cells. [J Immunol. 175: 5481-8.](#)
 10. Horrocks, P. *et al.* (2005) PfEMP1 expression is reduced on the surface of knobless *Plasmodium falciparum* infected erythrocytes. [J Cell Sci. 118: 2507-18.](#)
 11. Lozanoska-Ochser, B. *et al.* (2008) Expression of CD86 on human islet endothelial cells facilitates T cell adhesion and migration. [J Immunol. 181: 6109-16.](#)
 12. Norling, L.V. *et al.* (2008) Inhibitory control of endothelial galectin-1 on *in vitro* and *in vivo* lymphocyte trafficking. [FASEB J. 22: 682-90.](#)
 13. Baumer, Y. *et al.* (2011) Telomerase-based immortalization modifies the angiogenic/inflammatory responses of human coronary artery endothelial cells. [Exp Biol Med \(Maywood\). 236: 692-700.](#)
 14. Lask, A. *et al.* (2011) TCR-independent killing of B cell malignancies by anti-third-party CTLs: the critical role of MHC-CD8 engagement. [J Immunol. 187 \(4\): 2006-14.](#)
 15. Sommaggio, R. *et al.* (2012) Multiple Receptors Trigger Human NK Cell-Mediated Cytotoxicity against Porcine Chondrocytes. [J Immunol. 188: 2075-83.](#)
 16. Murphy, A.J. *et al.* (2013) Anti-inflammatory functions of apolipoprotein a-I and high-density lipoprotein are preserved in trimeric apolipoprotein a-I. [J Pharmacol Exp Ther. 344: 41-9.](#)
 17. Sumagin R *et al.* (2014) Transmigrated neutrophils in the intestinal lumen engage ICAM-1 to regulate the epithelial barrier and neutrophil recruitment. [Mucosal Immunol. 7 \(4\): 905-15.](#)
 18. Sugden SM *et al.* (2017) HIV-1 Vpu Downmodulates ICAM-1 Expression, Resulting in Decreased Killing of Infected CD4⁺ T Cells by NK Cells. [J Virol. 91 \(8\): pii: e02442-16.](#)
 19. Lennartz, F. *et al.* (2015) Mapping the Binding Site of a Cross-Reactive *Plasmodium falciparum* PfEMP1 Monoclonal Antibody Inhibitory of ICAM-1 Binding. [J Immunol. 195 \(7\): 3273-83.](#)

Storage Store at +4°C. DO NOT FREEZE.
 This product should be stored undiluted. This product is photosensitive and should be protected from light.
 Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Acknowledgements This product is provided under an intellectual property license from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchased product solely in research conducted by the buyer, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold

for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information Material Safety Datasheet documentation #20487 available at: 20487: <https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE-Alexa Fluor® 647 \(MCA928P647\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M378635:210222'

Printed on 18 Apr 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)