

## Datasheet: MCA1615PE

<b>Description:</b>	MOUSE ANTI HUMAN CD54:RPE
<b>Specificity:</b>	CD54
<b>Other names:</b>	ICAM-1
<b>Format:</b>	RPE
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	15.2
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

#### Target Species

Human

#### Species Cross Reactivity

Reacts with: Pig

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

#### Product Form

Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized

#### Reconstitution

Reconstitute with 1 ml distilled water

#### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
RPE 488nm laser	496	578

#### Buffer Solution

Phosphate buffered saline

<b>Preservative Stabilisers</b>	1% Bovine Serum Albumin 0.09% Sodium Azide 5% Sucrose
<b>Immunogen</b>	Human monocytes
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P05362</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">3383</a> ICAM1    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_322961
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse Sp2/0-Ag14 myeloma cell line
<b>Specificity</b>	<p><b>Mouse anti Human CD54 antibody, clone 15.2</b> recognizes the human CD54 cell surface antigen also known as intracellular Adhesion Molecule -1 (ICAM-1) or Major group rhinovirus receptor.</p> <p>CD54 is expressed by many cells following activation by inflammatory mediators. It is a 505 amino acid with an additional 27 amino acid signal peptide ~90 kDa single pass type I transmembrane glycoprotein bearing 5 Ig-like C2-type domains.</p> <p>Mouse anti Human CD54 antibody, clone 15.2 blocks CD54 function (<a href="#">Berendt et al. 1992</a>). Mouse anti Human CD54 antibody, clone 15.2 binds to an epitope on the N-terminal Ig-like domain within a region designated the L43 loop (<a href="#">Chakravorty and Craig 2005</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Dransfield, I. <i>et al.</i> (1992) Interaction of leukocyte integrins with ligand is necessary but not sufficient for function. <a href="#">J Cell Biol. 116:1527-35.</a></li> <li>2. Berendt, A. <i>et al.</i> (1992) The binding site on ICAM-1 for plasmodium falciparum-infected erythrocytes overlaps, but is distinct from, the LFA-1- binding site. <a href="#">Cell. 68: 71-81.</a></li> <li>3. Urquhart, P. <i>et al.</i> (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. <a href="#">J Pharmacol Exp Ther. 321 (2): 656-62.</a></li> <li>4. Baratin, M. <i>et al.</i> (2007) Dissection of the role of PfEMP1 and ICAM-1 in the sensing of <i>Plasmodium-falciparum</i>-infected erythrocytes by natural killer cells. <a href="#">PLoS One 2: e228.</a></li> <li>5. van Buul, J.D. <i>et al.</i> (2010) Inside-out regulation of ICAM-1 dynamics in TNF-alpha-activated endothelium. <a href="#">PLoS One 5: e11336.</a></li> <li>6. Diaz-Romero, J. <i>et al.</i> (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect bona fide dedifferentiation rather than amplification of progenitor cells. <a href="#">J Cell Physiol. 214: 75-83.</a></li> <li>7. Di Lorenzo, A. <i>et al.</i> (2011) Endothelial reticulon-4B (Nogo-B) regulates ICAM-1-mediated leukocyte transmigration and acute inflammation. <a href="#">Blood. 117: 2284-95.</a></li> <li>8. Porter, J.C. and Hall, A. (2009) Epithelial ICAM-1 and ICAM-2 regulate the egression of</li> </ol>

- human T cells across the bronchial epithelium. [FASEB J. 23: 492-502.](#)
9. Corvaisier, M. *et al.* (2005) V gamma 9V delta 2 T cell response to colon carcinoma cells. [J Immunol. 175: 5481-8.](#)
  10. Horrocks, P. *et al.* (2005) PfEMP1 expression is reduced on the surface of knobless *Plasmodium falciparum* infected erythrocytes. [J Cell Sci. 118: 2507-18.](#)
  11. Lozanoska-Ochser, B. *et al.* (2008) Expression of CD86 on human islet endothelial cells facilitates T cell adhesion and migration. [J Immunol. 181: 6109-16.](#)
  12. Norling, L.V. *et al.* (2008) Inhibitory control of endothelial galectin-1 on *in vitro* and *in vivo* lymphocyte trafficking. [FASEB J. 22: 682-90.](#)
  13. Baumer, Y. *et al.* (2011) Telomerase-based immortalization modifies the angiogenic/inflammatory responses of human coronary artery endothelial cells. [Exp Biol Med \(Maywood\). 236: 692-700.](#)
  14. Lask, A. *et al.* (2011) TCR-independent killing of B cell malignancies by anti-third-party CTLs: the critical role of MHC-CD8 engagement. [J Immunol. 187 \(4\): 2006-14.](#)
  15. Sommaggio, R. *et al.* (2012) Multiple Receptors Trigger Human NK Cell-Mediated Cytotoxicity against Porcine Chondrocytes. [J Immunol. 188: 2075-83.](#)
  16. Murphy, A.J. *et al.* (2013) Anti-inflammatory functions of apolipoprotein a-I and high-density lipoprotein are preserved in trimeric apolipoprotein a-I. [J Pharmacol Exp Ther. 344: 41-9.](#)
  17. Sumagin R *et al.* (2014) Transmigrated neutrophils in the intestinal lumen engage ICAM-1 to regulate the epithelial barrier and neutrophil recruitment. [Mucosal Immunol. 7 \(4\): 905-15.](#)
  18. Sugden SM *et al.* (2017) HIV-1 Vpu Downmodulates ICAM-1 Expression, Resulting in Decreased Killing of Infected CD4<sup>+</sup> T Cells by NK Cells. [J Virol. 91 \(8\): pii: e02442-16.](#)
  19. Lennartz, F. *et al.* (2015) Mapping the Binding Site of a Cross-Reactive *Plasmodium falciparum* PfEMP1 Monoclonal Antibody Inhibitory of ICAM-1 Binding. [J Immunol. 195 \(7\): 3273-83.](#)

---

**Storage** Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #20487 available at: 20487: <https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

## Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M375348:210104'

Printed on 18 Apr 2021

---

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)