

Datasheet: MCA1642F BATCH NUMBER 158675

| Description: | RAT ANTI HUMAN CD52:FITC |
|---------------|--------------------------|
| Specificity: | CD52 |
| Other names: | CAMPATH-1 |
| Format: | FITC |
| Product Type: | Monoclonal Antibody |
| Clone: | YTH34.5 |
| Isotype: | lgG2b |
| Quantity: | 0.1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | | | | Neat |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

or for

| Target Species | Human | | | |
|----------------|---------------------|---|------------------------|--------------------|
| Species Cross | Reacts with: Rhe | sus Monkey | | |
| Reactivity | reactivity is deriv | activity and working condit ed from testing within our l nications from the originaton. | laboratories, peer-rev | iewed publications |
| Product Form | Purified IgG conj | ugated to Fluorescein Isot | hiocyanate Isomer 1 | (FITC) - liquid |
| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) | |
| | FITC | 490 | 525 | |
| Preparation | Purified IgG prep | ared by affinity chromatog | raphy on Protein A fr | om tissue culture |

| Buffer Solution | Phosphate buffered saline |
|-----------------------------------|---|
| Preservative Stabilisers | 0.09% Sodium Azide 1% Bovine Serum Albumin |
| Approx. Protein Concentrations | IgG concentration 0.1 mg/ml |
| Immunogen | Human lymphocytes |
| External Database Links | UniProt: P31358 Related reagents |
| | Entrez Gene: 1043 CD52 Related reagents |
| Synonyms | CDW52, HE5 |
| RRID | AB_321471 |
| Specificity | Rat anti Human CD52 antibody, clone YTH34.5 recognizes the human CD52 antigen, also known as CAMPATH-1. The CD52 antigen is a remarkably small but heavily glycosylated peptide attached to the cell surface membrane via a GPI link (Xia et al. 1991). |
| | The apparent molecular mass of the native antigen on SDS-PAGE is 25-29 kDa, considerably reduced following N-glycanase treatment (Rowan et al. 1998). |
| | CD52 is expressed at high density by lymphocytes, monocytes, eosinophils, thymocytes and macrophages. It is expressed by most lymphoid derived malignancies, although expression on myeloma cells is variable. |
| | Humanized versions of CAMPATH-1 specific antibodies are currently in clinical trials for the treatment of a range of lymphoid malignancies (<u>Dearden et al. 2002</u> ; <u>Pettitt et al. 2012</u>). |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood. |
| References | Klangsinsirikul, P. et al. (2002) Campath-1G causes rapid depletion of circulating host dendritic cells (DCs) before allogeneic transplantation but does not delay donor DC reconstitution. Blood. 99: 2586-91. Ratzinger, G. et al. (2003) Differential CD52 expression by distinct myeloid dendritic cell subsets: implications for alemtuzumab activity at the level of antigen presentation in allogeneic graft-host interactions in transplantation. Blood. 101: 1422-9. Zand, M.S. et al. (2005) A renewable source of donor cells for repetitive monitoring of T- and B-cell alloreactivity. Am J Transplant. 5: 76-86. Westermann, J et al. (2005) CD52 Is Not a Promising Immunotherapy Target for Most |

- Patients with Multiple Myeloma International Journal of Hematology. 82 (3): 248-50.
- 5. Gopcsa, L. *et al.* (2005) Extensive flow cytometric characterization of plasmacytoid dendritic cell leukemia cells. Eur J Haematol. 75: 346-51.
- 6. Rodig SJ *et al.* (2006) Heterogeneous CD52 expression among hematologic neoplasms: implications for the use of alemtuzumab (CAMPATH-1H). <u>Clin Cancer Res. 12</u> (23): 7174-9.
- 7. Golay, J. *et al.* (2006) The sensitivity of acute lymphoblastic leukemia cells carrying the t(12;21) translocation to campath-1H-mediated cell lysis. <u>Haematologica</u>. 91: 322-30.
- 8. Miles, R.R. *et al.* (2007) Immunophenotypic identification of possible therapeutic targets in paediatric non-Hodgkin lymphomas: a children's oncology group report. <u>Br J Haematol.</u> 138: 506-12.
- 9. Chang, S.T. *et al.* (2007) CD52 expression in non-mycotic T- and NK/T-cell lymphomas. <u>Leuk Lymphoma</u>. 48: 117-21.
- 10. Piccaluga, P.P. *et al.* (2007) Expression of CD52 in peripheral T-cell lymphoma. Haematologica. 92: 566-7.
- 11. Reimer, P. *et al.* (2009) Autologous stem-cell transplantation as first-line therapy in peripheral T-cell lymphomas: results of a prospective multicenter study. <u>J Clin Oncol. 27:</u> 106-13.
- 12. Hu, Y. *et al.* (2009) Investigation of the mechanism of action of alemtuzumab in a human CD52 transgenic mouse model. <u>Immunology</u>. 128: 260-70.
- 13. Rizzo, K. *et al.* (2009) Novel CD19 expression in a peripheral T cell lymphoma: A flow cytometry case report with morphologic correlation. <u>Cytometry B Clin Cytom.</u> 76: 142-9.
- 14. Haniffa, M. *et al.* (2009) Differential rates of replacement of human dermal dendritic cells and macrophages during hematopoietic stem cell transplantation. <u>J Exp Med. 206:</u> 371-85.
- 15. Bisig, B. *et al.* (2013) CD30-positive peripheral T-cell lymphomas share molecular and phenotypic features. Haematologica. 98 (8): 1250-8.
- 16. Paulus, A. *et al.* (2015) Immunophenotyping of Waldenströms macroglobulinemia cell lines reveals distinct patterns of surface antigen expression: potential biological and therapeutic implications. <u>PLoS One. 10 (4): e0122338.</u>
- 17. Hotta, R. *et al.* (2016) CD52-Negative NK Cells Are Abundant in the Liver and Less Susceptible to Alemtuzumab Treatment. PLoS One. 11 (8): e0161618.
- 18. Buckstein, R. *et al.* (2016) Alemtuzumab and CHOP Chemotherapy for the Treatment of Aggressive Histology Peripheral T Cell Lymphomas: A Multi-Center Phase I Study. <u>Clin Lymphoma Myeloma Leuk.</u> 16 (1): 18-28.e4.
- 19. Craig, J.W. *et al.* (2018) Assessment of CD52 expression in "double-hit" and "double-expressor" lymphomas: Implications for clinical trial eligibility. <u>PLoS One. 13 (7):</u> e0199708.

Further Reading

- 1. Salisbury JR *et al.* (1994) Immunohistochemical analysis of CDw52 antigen expression in non-Hodgkin's lymphomas. <u>J Clin Pathol. 47 (4): 313-7.</u>
- 2. Hale G *et al.* (1998) Improving the outcome of bone marrow transplantation by using CD52 monoclonal antibodies to prevent graft-versus-host disease and graft rejection. Blood. 92 (12): 4581-90.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for

short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

| Guarantee | 12 months from date of despatch |
|----------------------------------|--|
| Health And Safety Information | Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1642F 10041 |
| Regulatory | For research purposes only |

Related Products

Recommended Negative Controls

RAT IgG2b NEGATIVE CONTROL:FITC (MCA6006F)

Email: antibody_sales_us@bio-rad.com

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

Fax: +1 919 878 3751

North & South Tel: +1 800 265 7376

America

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21

Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_uk@bio-rad.com Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385216:210513'

Printed on 01 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint