

Datasheet: MCA1651GA

Description:	MOUSE ANTI BOVINE CD205
Specificity:	CD205
Other names:	WC6 ANTIGEN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CC98
Isotype:	IgG2b
Quantity:	0.1 mg

Product Details

RRID AB_11203398

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Bovine

Species Cross Reactivity

Reacts with: Sheep
N.B. Antibody reactivity and working conditions may vary between species.

Product Form

Purified IgG - liquid

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.09% Sodium Azide

Carrier Free

Yes

Approx. Protein Concentrations

IgG concentration 1.0 mg/ml

Specificity

Mouse anti Bovine CD205 antibody, clone CC98 recognizes the bovine CD205 cell surface antigen, a ~210-220 kDa molecule expressed by T cells that are CD2+ve but not WC1+ve. CD205 is also expressed by B cells, and weakly stains B cell follicles.

Bovine CD205 has previously been described as the WC6 antigen ([Gliddon et al. 2004](#)).

Dendritic cells (veiled cells) in afferent lymph are strong expressors of CD205 as are dendritic cells in various other tissues.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Howard, C.J. & Naessens, J. (1993) Summary of workshop findings for cattle (tables 1 and 2). [Vet Immunol Immunopathol. 39 \(1-3\): 25-47.](#)
2. Howard, C.J. et al. (1996) Afferent lymph veiled cells stimulate proliferative responses in allogeneic CD4+ and CD8+ T cells but not gamma delta TCR+ T cells. [Immunology. 88 \(4\): 558-64.](#)
3. Naessens, J. et al. (1993) Cross-reactivity of workshop antibodies with cells from domestic and wild ruminants. [Vet Immunol Immunopathol. 39 \(1-3\): 283-90.](#)
4. Gliddon, D.R. et al. (2004) DEC-205 expression on migrating dendritic cells in afferent lymph. [Immunology. 111 \(3\): 262-72.](#)
5. Akesson, C.P. et al. (2008) Phenotypic characterisation of intestinal dendritic cells in sheep. [Dev Comp Immunol. 32: 837-49.](#)
6. Ferret-Bernard, S. et al. (2011) Mesenteric lymph node cells from neonates present a prominent IL-12 response to CpG oligodeoxynucleotide via an IL-15 feedback loop of amplification. [Vet Res. 42:19.](#)
7. Ferret-Bernard, S. et al. (2010) Cellular and molecular mechanisms underlying the strong neonatal IL-12 response of lamb mesenteric lymph node cells to R-848. [PLoS One. 5: e13705.](#)
8. Fach, S.J. et al. (2007) Neonatal ovine pulmonary dendritic cells support bovine respiratory syncytial virus replication with enhanced interleukin (IL)-4 And IL-10 gene transcripts. [Viral Immunol. 20: 119-30.](#)
9. Eicher, S.D. et al. (2011) β-Glucan plus ascorbic acid in neonatal calves modulates immune functions with and without *Salmonella enterica* serovar Dublin. [Vet Immunol Immunopathol. 142: 258-64.](#)
10. Olivier, M. et al. (2012) Capacities of Migrating CD1b Lymph Dendritic Cells to Present *Salmonella* Antigens to Naive T Cells [PLoS One. 7: e30430.](#)
11. Thonur, L. et al. (2012) Toll-like receptor gene expression in fresh and archived ovine pseudoafferent lymph DEC205+ dendritic cells. [J Comp Pathol. 147 \(2-3\): 296-304.](#)
12. Sigmundsdottir, H. et al. (2007) DCs metabolize sunlight-induced vitamin D3 to 'program' T cell attraction to the epidermal chemokine CCL27. [Nat Immunol. 8: 285-93.](#)
13. Fach, S.J. et al. (2007) Neonatal ovine pulmonary dendritic cells support bovine respiratory syncytial virus replication with enhanced interleukin (IL)-4 And IL-10 gene transcripts. [Viral Immunol. 20: 119-30.](#)
14. McNeilly, T.N. et al. (2006) Differential expression of cell surface markers by ovine respiratory tract dendritic cells. [J Histochem Cytochem. 54: 1021-30.](#)
15. Walters, A.A. et al. (2015) Assessment of the enhancement of PLGA nanoparticle uptake by dendritic cells through the addition of natural receptor ligands and monoclonal antibody. [Vaccine. 33 \(48\): 6588-95.](#)
16. Lund, H. et al. (2016) Transient Migration of Large Numbers of CD14(++) CD16(+) Monocytes to the Draining Lymph Node after Onset of Inflammation. [Front Immunol. 7: 322.](#)
17. Uhde, A-K. et al. (2017) Evaluation of a panel of antibodies for the immunohistochemical

identification of immune cells in paraffin-embedded lymphoid tissues of new- and old-world camelids [Vet Immunol Immunopathol. Jan 17 \[Epub ahead of print\]](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Human Anti Mouse IgG2b (HCA038...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®680](#),
[DyLight®800](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL \(MCA691\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M334393:181127'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)