

## Datasheet: MCA1654A488

<b>Description:</b>	MOUSE ANTI BOVINE CD8 BETA:Alexa Fluor® 488
<b>Specificity:</b>	CD8 BETA
<b>Format:</b>	ALEXA FLUOR® 488
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	CC58
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS/1ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**Target Species** Bovine

**Species Cross Reactivity** Reacts with: Sheep, Goat, Water Buffalo  
**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

**Product Form** Purified IgG conjugated to Alexa Fluor® 488- liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®488	495	519

**Preparation** Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

**Buffer Solution** Phosphate buffered saline

<b>Preservative</b>	0.09% sodium azide (NaN <sub>3</sub> )
<b>Stabilisers</b>	1% bovine serum albumin
<b>Approx. Protein Concentrations</b>	Ig concentration 0.05 mg/ml
<b>Immunogen</b>	Bovine leucocytes
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">A7YW30</a> <a href="#">Related reagents</a>
<b>Specificity</b>	<p><b>Mouse anti Bovine CD8 beta antibody, clone CC58</b> recognizes an epitope associated with the bovine CD8 beta chain.</p> <p>CD8 is usually expressed as an α/β heterodimer. The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. The CD8 antigen, acting as a coreceptor, and the T-cell receptor on the T lymphocyte recognize antigens displayed by an antigen presenting cell (APC) in the context of class I MHC molecules.</p> <p>Mouse anti Bovine CD8 beta antibody, clone CC58 has been successfully used for the immunohistochemical detection of CD8 on formalin fixed, paraffin embedded placental tissue from water buffalo (<a href="#">Cantón <i>et al.</i> 2014</a>).</p>
<b>Flow Cytometry</b>	Use 10µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl
<b>References</b>	<ol style="list-style-type: none"> <li>1. Suraud, V. <i>et al.</i> (2008) Acute infection by conjunctival route with <i>Brucella melitensis</i> induces IgG+ cells and IFN-gamma producing cells in peripheral and mucosal lymph nodes in sheep. <a href="#">Microbes Infect. 10: 1370-8.</a></li> <li>2. Howard, C.J. &amp; Naessens, J. (1993) Summary of workshop findings for cattle (tables 1 and 2). <a href="#">Vet Immunol Immunopathol. 39 (1-3): 25-47.</a></li> <li>3. Naessens, J. <i>et al.</i> (1997) Nomenclature and characterization of leukocyte differentiation antigens in ruminants. <a href="#">Immunol Today. 18 (8): 365-8.</a></li> <li>4. Hein, W.R. <i>et al.</i> (1991) Summary of workshop findings for leukocyte antigens of sheep. <a href="#">Vet Immunol Immunopathol. 27 (1-3): 28-30.</a></li> <li>5. Gerner, W. <i>et al.</i> (2009) Identification of major histocompatibility complex restriction and anchor residues of foot-and-mouth disease virus-derived bovine T-cell epitopes. <a href="#">J Virol. 83: 4039-50.</a></li> <li>6. Gerner, W. <i>et al.</i> (2010) Sensitive detection of Foxp3 expression in bovine lymphocytes by flow cytometry. <a href="#">Vet Immunol Immunopathol. 138: 154-8.</a></li> <li>7. MacHugh, N.D. and Sopp, P. (1991) Individual antigens of cattle. Bovine CD8 (BoCD8). <a href="#">Vet Immunol Immunopathol. 27: 65-9.</a></li> <li>8. Soltys, J. and Quinn, M.T. (1999) Selective recruitment of T-cell subsets to the udder during staphylococcal and streptococcal mastitis: analysis of lymphocyte subsets and adhesion molecule expression. <a href="#">Infect Immun. 67: 6293-302.</a></li> <li>9. Cantón, G.J. <i>et al.</i> (2014) Characterization of immune cell infiltration in the placentome of water buffaloes (<i>Bubalus bubalis</i>) infected with <i>neospora caninum</i> during pregnancy. <a href="#">J</a></li> </ol>

[Comp Pathol. 150: 463-8.](#)

10. Wattegedera, S.R. *et al.* (2017) Enhancing the toolbox to study IL-17A in cattle and sheep. [Vet Res. 48 \(1\): 20.](#)

11. Hecker, Y.P. *et al.* (2015) Cell mediated immune responses in the placenta following challenge of vaccinated pregnant heifers with *Neospora caninum*. [Vet Parasitol. 214 \(3-4\): 247-54.](#)

12. Okino, C.H. *et al.* (2020) A polymorphic CD4 epitope related to increased susceptibility to *Babesia bovis*. in Canchim calves. [Vet Immunol Immunopathol. 230: 110132.](#)

---

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1654A488>  
10041

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 488 \(MCA928A488\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M410963:221031'

**Printed on 18 Jan 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)