

## Datasheet: MCA1655G

<b>Description:</b>	MOUSE ANTI BOVINE WC1
<b>Specificity:</b>	WC1
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	CC101
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.25 mg

## Product Details

**RRID** AB\_324174

**Applications** This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Functional Assays			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**Target Species** Bovine

**Species Cross Reactivity** Reacts with: Pig, Sheep  
**N.B.** Antibody reactivity and working conditions may vary between species.

**Product Form** Purified IgG - liquid

**Preparation** Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

**Buffer Solution** Phosphate buffered saline

**Preservative Stabilisers** 0.09% Sodium Azide

**Approx. Protein Concentrations** IgG concentration 1 mg/ml

**Immunogen**

Con A stimulated bovine lymphocytes

**External Database Links****UniProt:**[P30205](#) [Related reagents](#)**Entrez Gene:**[338056](#) CD163L1 [Related reagents](#)**Specificity**

**Mouse anti Bovine WC1 antibody, clone CC101**, recognizes a subset of WC1<sup>+</sup> T-cells expressing the WC1.1 isoform ([MacHugh et al. 1993](#)). The bovine WC1 cell surface antigen is expressed by a population of gamma/delta T-cells that lack CD2, CD4 and CD8, but express CD3. WC1 expression appears to be heterogeneous and antibodies to this cluster show differing reaction patterns ([Crocker et al. 1993](#)).

Mouse anti bovine WC1, clone CC101, immunoprecipitates a 215 kDa molecule from bovine cells and also recognizes the swine homolog of WC1, which is a 180 kDa molecule. In pigs, the 180 kDa molecule is expressed by a gamma/delta TCR positive T-cell population that also lack CD2, CD4 and CD8 ([Carr et al. 1994](#)).

**Flow Cytometry**Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul.**References**

1. Carr, M.M. *et al.* (1994) Expression on porcine gamma/delta lymphocytes of a phylogenetically conserved surface antigen previously restricted in expression to ruminant gamma/delta T lymphocytes. [Immunology 81: 36-40.](#)
2. Howard, C.J. & Naessens, J. (1993) Summary of workshop findings for cattle. [Vet Immunol Immunopathol. 39: 25-48.](#)
3. MacHugh, N. *et al.* (1993) Clustering of monoclonal antibodies recognizing different members of the WC1 gene family. [Vet Immunol Immunopathol. 39: 155-60.](#)
4. Crocker, G. *et al.* (1993) Analysis of the gamma/delta T cell restricted antigen WC1. [Vet Immunol Immunopathol. 39: 137-44.](#)
5. Lund, B. *et al.* (1993) Expression of T19 (WC1) molecules by ovine lymphocytes. [Vet Immunol Immunopathol. 39: 145-53.](#)
6. Schröder, A.C. & Hamann, J. (2005) The influence of technical factors on differential cell count in milk. [J Dairy Res. 72: 153-8.](#)
7. Liu, X. *et al.* (2014) Crusted scabies is associated with increased IL-17 secretion by skin T cells. [Parasite Immunol. 36: 594-604.](#)
8. Patarroyo, J.H. *et al.* (2009) Immune response of bovines stimulated by synthetic vaccine SBm7462 against *Rhipicephalus (Boophilus) microplus*. [Vet Parasitol. 166: 333-9.](#)
9. Heiser, A. *et al.* (2015) Grazing dairy cows had decreased interferon- $\gamma$ , tumor necrosis factor, and interleukin-17, and increased expression of interleukin-10 during the first week after calving. [J Dairy Sci. 98: 937-46.](#)
10. Takamatsu, H.H. *et al.* (2006) Porcine gammadelta T cells: possible roles on the innate and adaptive immune responses following virus infection. [Vet Immunol Immunopathol. 112: 49-61.](#)
11. Sedlak, C. *et al.* (2014) IL-12 and IL-18 induce interferon- $\gamma$  production and de novo CD2 expression in porcine  $\gamma\delta$  T cells. [Dev Comp Immunol. 47: 115-22.](#)
12. Herry, V. *et al.* (2017) Local immunization impacts the response of dairy cows to *Escherichia coli* mastitis. [Sci Rep. 7 \(1\): 3441.](#)

**Further Reading**

1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)
2. Wijngaard, P. *et al.* (1992) Molecular characterization of the WC1 antigen expressed specifically on bovine CD4-CD8- gamma delta T lymphocytes. [J Immunol. 149: 3273-7.](#)

<b>Storage</b>	Store at +4°C or at -20°C if preferred.  This product should be stored undiluted.  Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
<b>Shelf Life</b>	18 months from date of despatch.
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: 10040: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR8...)	<a href="#">DyLight@800</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Human Anti Mouse IgG2a (HCA037...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight@488</a> , <a href="#">DyLight@549</a> , <a href="#">DyLight@649</a> , <a href="#">DyLight@680</a> , <a href="#">DyLight@800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M334403:181127'

Printed on 02 Jan 2019