

Datasheet: MCA1655GA

BATCH NUMBER 149995

Description:	MOUSE ANTI BOVINE WC1
Specificity:	WC1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CC101
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Functional Assays			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Bovine

Species Cross Reactivity

Reacts with: Pig, Sheep

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture

supernatant

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.09% Sodium Azide (NaN₃)

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1.0 mg/ml

Immunogen Con A stimulated bovine lymphocytes

External Database Links

UniProt:

[P30205](#)

[Related reagents](#)

Entrez Gene:

[338056](#)

CD163L1

[Related reagents](#)

Specificity

Mouse anti Bovine WC1 antibody, clone CC101, recognizes a subset of WC1⁺ T-cells expressing the WC1.1 isoform (MacHugh *et al.* 1993). The bovine WC1 cell surface antigen is expressed by a population of gamma/delta T-cells that lack CD2, CD4 and CD8, but express CD3. WC1 expression appears to be heterogeneous and antibodies to this cluster show differing reaction patterns ([Crocker *et al.* 1993](#)).

Mouse anti bovine WC1, clone CC101, immunoprecipitates a 215 kDa molecule from bovine cells and also recognizes the swine homolog of WC1, which is a 180 kDa molecule. In pigs, the 180 kDa molecule is expressed by a gamma/delta TCR positive T-cell population that also lack CD2, CD4 and CD8 ([Carr *et al.* 1994](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Carr, M.M. *et al.* (1994) Expression on porcine gamma/delta lymphocytes of a phylogenetically conserved surface antigen previously restricted in expression to ruminant gamma/delta T lymphocytes. [Immunology 81: 36-40.](#)
2. Howard, C.J. & Naessens, J. (1993) Summary of workshop findings for cattle. [Vet Immunol Immunopathol. 39: 25-48.](#)
3. MacHugh, N. *et al.* (1993) Clustering of monoclonal antibodies recognizing different members of the WC1 gene family. [Vet Immunol Immunopathol. 39: 155-60.](#)
4. Crocker, G. *et al.* (1993) Analysis of the gamma/delta T cell restricted antigen WC1. [Vet Immunol Immunopathol. 39: 137-44.](#)
5. Lund, B. *et al.* (1993) Expression of T19 (WC1) molecules by ovine lymphocytes. [Vet Immunol Immunopathol. 39: 145-53.](#)
6. Schröder, A.C. & Hamann, J. (2005) The influence of technical factors on differential cell count in milk. [J Dairy Res. 72: 153-8.](#)
7. Liu, X. *et al.* (2014) Crusted scabies is associated with increased IL-17 secretion by skin T cells. [Parasite Immunol. 36: 594-604.](#)

8. Patarroyo, J.H. *et al.* (2009) Immune response of bovines stimulated by synthetic vaccine SBm7462 against *Rhipicephalus (Boophilus) microplus*. [Vet Parasitol. 166: 333-9.](#)
9. Heiser, A. *et al.* (2015) Grazing dairy cows had decreased interferon- γ , tumor necrosis factor, and interleukin-17, and increased expression of interleukin-10 during the first week after calving. [J Dairy Sci. 98: 937-46.](#)
10. Takamatsu, H.H. *et al.* (2006) Porcine gammadelta T cells: possible roles on the innate and adaptive immune responses following virus infection. [Vet Immunol Immunopathol. 112: 49-61.](#)
11. Sedlak, C. *et al.* (2014) IL-12 and IL-18 induce interferon- γ production and de novo CD2 expression in porcine $\gamma\delta$ T cells. [Dev Comp Immunol. 47: 115-22.](#)
12. Herry, V. *et al.* (2017) Local immunization impacts the response of dairy cows to *Escherichia coli* mastitis. [Sci Rep. 7 \(1\): 3441.](#)
13. Hussen, J. *et al.* (2018) Expression Patterns of Cell Adhesion Molecules on CD4+ T Cells and WC1+ T Cells in the Peripheral Blood of Dromedary Camels. [Pakistan Veterinary Journal. 38 \(03\): 231-236.](#)

Further Reading	<ol style="list-style-type: none"> 1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. Vet Res. 39: 54. 2. Wijngaard, P. <i>et al.</i> (1992) Molecular characterization of the WC1 antigen expressed specifically on bovine CD4-CD8- gamma delta T lymphocytes. J Immunol. 149: 3273-7.
------------------------	--

Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
----------------	--

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	<p>Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1655GA</p> <p>10040</p>
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@550 , DyLight@650 , DyLight@680 , DyLight@800 ,

[FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR9...)

[FITC](#)

Goat Anti Mouse IgG (STAR77...)

[HRP](#)

Goat Anti Mouse IgG (Fc) (STAR120...)

[FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M365584:200529'

Printed on 24 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)