

Datasheet: MCA1682GT

BATCH NUMBER 162219

Description:	MOUSE ANTI HUMAN BRCA1 (N-TERMINAL)
Specificity:	BRCA1 (N-TERMINAL)
Other names:	BREAST CANCER TYPE 1 SUSCEPTIBILITY PROTEIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	MS110
Isotype:	IgG1
Quantity:	50 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin (2)	▪			
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			1.0ug/ml
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) **This product requires antigen retrieval using heat treatment prior to staining of frozen sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.** See [Yoshikawa, K.et al.](#) for details.

(2) **This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.** See [Yoshikawa, K.et al.](#) for details.

Target Species	Human
-----------------------	-------

Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	TRIS-glycine buffered saline, NaCl
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Recombinant protein corresponding to the N-Terminal region of human BRCA1.
External Database Links	<p>UniProt: P38398 Related reagents</p> <p>Entrez Gene: 672 BRCA1 Related reagents</p>
Synonyms	RNF53
Fusion Partners	Spleen cells from immunized mice were fused with cells of the mouse NSI myeloma cell line.
Specificity	<p>Mouse anti Human BRCA1 antibody, clone MS110 recognizes an epitope within the 304 amino acid N-Terminal (NT) region of human BRCA1, otherwise known as Breast cancer type 1 susceptibility protein, a tumor suppressor gene and major player in DNA damage repair, predominantly expressed in the nucleus during the S/G2 phase of the cell cycle.</p> <p>Deleterious BRCA1 mutations corralate with increased risk of cancers in both males and females including breast, pancreatic and prostate cancer, although in males pancreatic and prostate cancer appear to be more strongly associated with BRCA2 gene mutations.</p> <p>BRCA1 is a key marker of triple-negative breast cancer/TNBC (ER-/PR-/HER2-), a high risk aggressive cancer which makes up about 15% of invasive breast cancers, and which lacks the benefit of specific therapy that targets the three major proteins ER/PR/HER2. Triple-negative tumors are predominantly basal-like, poorly differentiated and of higher histological grade. Younger women have an increased rate of basal or BRCA related TNBC, compared with the higher proportion of apocrine, normal-like and rare subtypes of TNBC, seen in older women.</p> <p>Mouse anti human BRCA1 antibody, clone MS110 is suitable for use in both immunohistochemical and immunofluorescence staining of human breast and ovarian tissue, and clone MS110 is widely used and reported (Milner <i>et al.</i> 2013).</p>

Histology Positive Control Tissue	Breast carcinoma
Western Blotting	Mouse anti Human BRCA1 detects a band of approximately 220kDa in HeLa nuclear extract. A smaller band of 85kDa may also be seen, which may represent post-translational modification.
References	<ol style="list-style-type: none"> 1. Scully, R. <i>et al.</i> (1996) Location of BRCA1 in human breast and ovarian cancer cells. Science. 272 (5258): 123-6. 2. Yoshikawa, K. <i>et al.</i> (1999) Reduction of BRCA1 protein expression in Japanese sporadic breast carcinomas and its frequent loss in BRCA1-associated cases. Clin Cancer Res. 5 (6): 1249-61. 3. Scully, R. <i>et al.</i> (1997) Association of BRCA1 with Rad51 in mitotic and meiotic cells. Cell. 88: 265-75. 4. Chen, J. (2000) Ataxia telangiectasia-related protein is involved in the phosphorylation of BRCA1 following deoxyribonucleic acid damage. Cancer Res. 60: 5037-9. 5. Fraser, J.A. <i>et al.</i> (2003) A role for BRCA1 in sporadic breast cancer. Br J Cancer. 88: 1263-70. 6. Kleiman, F.E. <i>et al.</i> (2005) BRCA1/BARD1 inhibition of mRNA 3' processing involves targeted degradation of RNA polymerase II. Genes Dev. 19: 1227-37. 7. Pageau, G.J. and Lawrence, J.B. (2006) BRCA1 foci in normal S-phase nuclei are linked to interphase centromeres and replication of pericentric heterochromatin. J Cell Biol. 175: 693-701.
Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10511 available at: https://www.bio-rad-antibodies.com/SDS/MCA1682GT10511
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP

Goat Anti Mouse IgG (STAR70...)

[FITC](#)

Goat Anti Mouse IgG (H/L) (STAR117...)

[Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR9...)

[FITC](#)

Goat Anti Mouse IgG (Fc) (STAR120...)

[FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M389321:210806'

Printed on 11 Sep 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)