

Datasheet: MCA1682GT

## **BATCH NUMBER 170321**

Description:MOUSE ANTI HUMAN BRCA1 (N-TERMINAL)Specificity:BRCA1 (N-TERMINAL)Other names:BREAST CANCER TYPE 1 SUSCEPTIBILITY PROTEINFormat:PurifiedProduct Type:Monoclonal AntibodyClone:MS110Isotype:IgG1Quantity:50 μg		
Other names: BREAST CANCER TYPE 1 SUSCEPTIBILITY PROTEIN  Format: Purified  Product Type: Monoclonal Antibody  Clone: MS110  Isotype: IgG1	Description:	MOUSE ANTI HUMAN BRCA1 (N-TERMINAL)
Format: Purified  Product Type: Monoclonal Antibody  Clone: MS110  Isotype: IgG1	Specificity:	BRCA1 (N-TERMINAL)
Product Type: Monoclonal Antibody Clone: MS110 Isotype: IgG1	Other names:	BREAST CANCER TYPE 1 SUSCEPTIBILITY PROTEIN
Clone: MS110 Isotype: IgG1	Format:	Purified
Isotype: IgG1	Product Type:	Monoclonal Antibody
· · · · · · · · · · · · · · · · · · ·	Clone:	MS110
Quantity: 50 μg	Isotype:	lgG1
	Quantity:	50 μg

# **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			
Immunohistology - Frozen (1)	•			
Immunohistology - Paraffin (2)	•			
ELISA			•	
Immunoprecipitation	•			
Western Blotting	•			1.0ug/ml
Immunofluorescence	•			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

- (1) This product requires antigen retrieval using heat treatment prior to staining of frozen sections. Sodium citrate buffer pH 6.0 is recommended for this purpose. See <u>Yoshikawa, K.et al.</u> for details.
- (2) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose. See Yoshikawa, K.et al. for details.

Target Species	Human

Product Form	Purified IgG - liquid		
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	TRIS-glycine buffered saline, NaCl		
Preservative Stabilisers	0.05% Sodium Azide		
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml		
Immunogen	Recombinant protein corresponding to the N-Terminal region of human BRCA1.		
External Database Links	UniProt:  P38398 Related reagents  Entrez Gene:  672 BRCA1 Related reagents		
Synonyms	RNF53		
Fusion Partners	Spleen cells from immunised mice were fused with cells of the mouse NSI myeloma cell line.		

## **Specificity**

**Mouse anti Human BRCA1 antibody, clone MS110** recognizes an epitope within the 304 amino acid N-Terminal (NT) region of human BRCA1, otherwise known as Breast cancer type 1 susceptibility protein, a tumor suppressor gene and major player in DNA damage repair, predominantly expressed in the nucleus during the S/G2 phase of the cell cycle.

Along with BRCA2, BRCA1 is a high risk gene which is associated with hereditary breast and ovarian cancers, particularly at a younger age of diagnosis. Women carrying the BRCA1 mutation have a 50-95% chance of developing breast cancer in later life, but genetic screening and increased awareness of preventative surgery, can reduce this risk significantly. Deleterious BRCA1 mutations may also increase the risk of other cancers in both males and females including pancreatic cancer, although in males pancreatic and prostate cancer appear to be more strongly associated with BRCA2 gene mutations.

BRCA1 is a key marker of triple-negative breast cancer/TNBC (ER-/PR-/HER2-), a high risk aggressive cancer which makes up about 15% of invasive breast cancers, and which lacks the benefit of specific therapy that targets the three major proteins ER/PR/HER2. Triple-negative tumors are predominantly basal-like, poorly differentiated and of higher <a href="histological grade">histological grade</a>. Younger women have an increased rate of basal or BRCA related TNBC, compared with the higher proportion of apocrine, normal-like and rare subtypes of TNBC, seen in older women.

	tissue, and clone MS110 is a widely recognized antibody for use in cancer studies (Milner et al 2013).
Histology Positive Control Tissue	Breast carcinoma
Western Blotting	Mouse anti Human BRCA1 detects a band of approximately 220kDa in HeLa nuclear extract. A smaller band of 85kDa may also be seen, which may represent post-translational modification.
References	<ol> <li>Scully, R. et al. (1996) Location of BRCA1 in human breast and ovarian cancer cells. Science. 272 (5258): 123-6.</li> <li>Yoshikawa, K. et al. (1999) Reduction of BRCA1 protein expression in Japanese sporadic breast carcinomas and its frequent loss in BRCA1-associated cases. Clin Cancer Res. 5 (6): 1249-61.</li> <li>Scully, R. et al. (1997) Association of BRCA1 with Rad51 in mitotic and meiotic cells. Cell. 88: 265-75.</li> <li>Chen, J. (2000) Ataxia telangiectasia-related protein is involved in the phosphorylation of BRCA1 following deoxyribonucleic acid damage. Cancer Res. 60: 5037-9.</li> <li>Fraser, J.A. et al. (2003) A role for BRCA1 in sporadic breast cancer. Br J Cancer. 88: 1263-70.</li> <li>Kleiman, F.E. et al. (2005) BRCA1/BARD1 inhibition of mRNA 3' processing involves targeted degradation of RNA polymerase II. Genes Dev. 19: 1227-37.</li> <li>Pageau, G.J. and Lawrence, J.B. (2006) BRCA1 foci in normal S-phase nuclei are linked to interphase centromeres and replication of pericentric heterochromatin. J Cell Biol. 175: 693-701.</li> </ol>
Storage	Store at +4°C or at -20°C if preferred.  This product should be stored undiluted.  Storage in frost free freezers is not recommended Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10511 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1682GT">https://www.bio-rad-antibodies.com/SDS/MCA1682GT</a> 10511
Regulatory	For research purposes only

Mouse anti human BRCA1 antibody, clone MS110 is suitable for use in both

immunohistochemical and immunofluorescence staining of human breast and ovarian

# **Related Products**

**Recommended Secondary Antibodies** 

Rabbit Anti Mouse IgG (STAR12...)

Goat Anti Mouse IgG IgA IgM (STAR87...)

HRP

Goat Anti Mouse IgG (STAR76...)

RPE

Rabbit Anti Mouse IgG (STAR13...)

HRP

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) <u>FITC</u>
Goat Anti Mouse IgG (STAR77...) <u>HRP</u>

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

## **Recommended Negative Controls**

## MOUSE IgG1 NEGATIVE CONTROL (MCA928)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365593:200529'

### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint