

Datasheet: MCA1701

BATCH NUMBER 155531

Description:	MOUSE ANTI p53 (aa20-25)
Specificity:	p53 (aa20-25)
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	DO-1
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			1/25
ELISA	▪			
Immunoprecipitation	▪			
Western Blotting	▪			1/1000

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species

Human

Species Cross Reactivity

Reacts with: Bovine, Cat, Horse, Green Monkey

Does not react with: Mouse, Rat

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Recombinant human p53.
External Database Links	<p>UniProt: P04637 Related reagents</p> <p>Entrez Gene: 7157 TP53 Related reagents</p>
Synonyms	P53
RRID	AB_322633
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse X63Ag8.653 myeloma cell line.
Specificity	<p>Mouse anti Human p53 antibody, clone DO-1 recognizes the human p53 tumor suppressor protein, also known as cellular tumor antigen p53 or NY-CO-13. Clone DO-1 binds to both wild type and mutant forms of the p53 protein found in various malignancies (Kern et al. 1992). p53 is important in multicellular organisms, where it regulates cell cycle progression to allow DNA repair or apoptosis in the case of irreparably damaged cells (Haupt et al. 2003) and thus functions as a tumor suppressor that is involved in preventing cancer. Mutations in the p53 gene are found in about half the cases of human cancer (Joerger and Fersht 2007)</p> <p>Mouse anti Human p53 antibody, clone DO-1 recognizes an epitope at the N-terminal end of p53 between amino acids 20-25, common to isoforms 1-3 of p53.</p>
Histology Positive Control Tissue	Colon or breast carcinoma
References	<ol style="list-style-type: none"> Vojtěšek B <i>et al.</i> (1992) An immunochemical analysis of the human nuclear phosphoprotein p53. New monoclonal antibodies and epitope mapping using recombinant p53. J Immunol Methods. 151 (1-2): 237-44. Levesque, M.A. <i>et al.</i> (1995) Time-resolved immunofluorometric assay of p53 protein.

[Clin Chem. 41 \(12 Pt 1\): 1720-9.](#)

3. Carvalho, T. *et al.* (2009) Immunohistochemical evaluation of vascular urinary bladder tumors from cows with enzootic hematuria. [Vet Pathol. 46 \(2\): 211-21.](#)

4. Phillips, A. *et al.* (2010) HDMX-L is expressed from a functional p53-responsive promoter in the first intron of the HDMX gene and participates in an autoregulatory feedback loop to control p53 activity. [J Biol Chem. 285 \(38\): 29111-27.](#)

5. Bergman, L.M. *et al.* (2009) CtBPs promote cell survival through the maintenance of mitotic fidelity. [Mol Cell Biol. 29: 4539-51.](#)

6. Hietanen, S. *et al.* (2000) Activation of p53 in cervical carcinoma cells by small molecules. [Proc Natl Acad Sci U S A. 97 \(15\): 8501-6.](#)

7. Phelps, M. *et al.* (2003) p53-independent activation of the hdm2-P2 promoter through multiple transcription factor response elements results in elevated hdm2 expression in estrogen receptor alpha-positive breast cancer cells. [Cancer Res. 63: 2616-23.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1701>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)

Goat Anti Mouse IgG (STAR76...) [RPE](#)

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Goat Anti Mouse IgG (STAR70...) [FITC](#)

Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Goat Anti Mouse IgG (STAR77...) [HRP](#)

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M365598:200529'

Printed on 18 Jan 2024
