

Datasheet: MCA1701B

BATCH NUMBER 0912R

Description:	MOUSE ANTI p53 (aa20-25):Biotin
Specificity:	p53 (aa20-25)
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	DO-1
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Paraffin	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Bovine, Cat, Horse, Green Monkey</p> <p>Does not react with: Mouse, Rat</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG conjugated to Biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	<p>0.09% Sodium Azide</p> <p>1% Bovine Serum Albumin</p>

Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Recombinant human p53.
External Database Links	<p>UniProt: P04637 Related reagents</p> <p>Entrez Gene: 7157 TP53 Related reagents</p>
Synonyms	P53
RRID	AB_323761
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse X63Ag8.653 myeloma cell line.
Specificity	<p>Mouse anti Human p53 antibody, clone DO-1 recognizes the human p53 tumor suppressor protein, also known as cellular tumor antigen p53 or NY-CO-13. Clone DO-1 binds to both wild type and mutant forms of the p53 protein found in various malignancies (Kern et al. 1992). p53 is important in multicellular organisms, where it regulates cell cycle progression to allow DNA repair or apoptosis in the case of irreparably damaged cells (Haupt et al. 2003) and thus functions as a tumor suppressor that is involved in preventing cancer. Mutations in the p53 gene are found in about half the cases of human cancer (Joerger and Fersht 2007)</p> <p>Mouse anti Human p53 antibody, clone DO-1 recognizes an epitope at the N-terminal end of p53 between amino acids 20-25, common to isoforms 1-3 of p53.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
References	<ol style="list-style-type: none"> 1. Vojtěšek B <i>et al.</i> (1992) An immunochemical analysis of the human nuclear phosphoprotein p53. New monoclonal antibodies and epitope mapping using recombinant p53. J Immunol Methods. 151 (1-2): 237-44. 2. Levesque, M.A. <i>et al.</i> (1995) Time-resolved immunofluorometric assay of p53 protein. Clin Chem. 41 (12 Pt 1): 1720-9. 3. Carvalho, T. <i>et al.</i> (2009) Immunohistochemical evaluation of vascular urinary bladder tumors from cows with enzootic hematuria. Vet Pathol. 46 (2): 211-21. 4. Phillips, A. <i>et al.</i> (2010) HDMX-L is expressed from a functional p53-responsive promoter in the first intron of the HDMX gene and participates in an autoregulatory feedback loop to control p53 activity. J Biol Chem. 285 (38): 29111-27. 5. Bergman, L.M. <i>et al.</i> (2009) CtBPs promote cell survival through the maintenance of mitotic fidelity. Mol Cell Biol. 29: 4539-51. 6. Hietanen, S. <i>et al.</i> (2000) Activation of p53 in cervical carcinoma cells by small molecules. Proc Natl Acad Sci U S A. 97 (15): 8501-6. 7. Phelps, M. <i>et al.</i> (2003) p53-independent activation of the hdm2-P2 promoter through

multiple transcription factor response elements results in elevated hdm2 expression in estrogen receptor alpha-positive breast cancer cells. [Cancer Res. 63: 2616-23.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1701B>

Regulatory

For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M365599:200529'

Printed on 09 Jul 2025