

## Datasheet: MCA1703

**BATCH NUMBER 154721**

<b>Description:</b>	MOUSE ANTI p53 (aa20-25)
<b>Specificity:</b>	p53 (aa20-25)
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	DO-7
<b>Isotype:</b>	IgG2b
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (2)	▪			1/1000
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			1/1000

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1)Membrane permeabilization is required for this application. Bio-Rad recommend the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.**

**(2)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.Sodium citrate buffer pH 6.0 is recommended for this purpose.**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	<p>Reacts with: Bovine</p> <p>Does not react with:Mouse, Rat</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for</p>

further information.

<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Recombinant human p53.
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P04637</a> <a href="#">Related reagents</a>  <b>Entrez Gene:</b> <a href="#">7157</a> TP53 <a href="#">Related reagents</a>
<b>Synonyms</b>	P53
<b>RRID</b>	AB_322635
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63Ag8.653 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti p53 antibody, clone DO-7</b> recognizes the human 53 kDa p53 tumour suppressor protein, also known as Cellular tumor antigen p53 or Antigen NY-CO-13, encoded by the TP53 gene. p53 is a 393 amino acid protein with an N-terminal transactivation domain, followed by a proline-rich region and a DNA binding domain in the central core region. The C-terminal region contains a tetramerization domain and a terminal regulatory domain (<a href="#">Joerger et al. 2010</a>).</p> <p>p53 is intimately involved in a number of signaling pathways controlling cell division, cycling and apoptosis (<a href="#">Haupt et al. 2003</a>) and is thus a potent cancer suppressor. In normal cells the level of p53 expression is low but can be induced by DNA damage or other stress signals (<a href="#">Takagi et al. 2005</a>). Activation of p53 leads to growth arrest through its interaction with p21, GADD45 and 14-3-3<math>\sigma</math>, DNA repair and potentially apoptosis through interaction with Bax, Apaf-1, PUMA and NoxA (<a href="#">Thakur et al. 2010</a>). p53 is critically regulated by Mdm2 which can trigger p53 degradation by a ubiquitin dependent system (<a href="#">Moll and Petrenko 2003</a>)</p> <p>Mouse anti p53 antibody, clone DO-7 recognizes an epitope at the N-terminal end of p53</p>

between amino acids 20-25, binding to both wild type and mutant forms. Clone DO-7 is not expected to recognize the multiple [isoforms](#) lacking the N-terminal region.

<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul.
<b>Histology Positive Control Tissue</b>	Colon or breast carcinoma
<b>References</b>	<ol style="list-style-type: none"><li>1. Vojtěšek, B. <i>et al.</i> (1992) An immunochemical analysis of the human nuclear phosphoprotein p53. New monoclonal antibodies and epitope mapping using recombinant p53. <a href="#">J Immunol Methods. 151 (1-2): 237-44.</a></li><li>2. Xinarianos, G. <i>et al.</i> (2002) p53 status correlates with the differential expression of the DNA mismatch repair protein MSH2 in non-small cell lung carcinoma. <a href="#">Int J Cancer. 101: 248-52.</a></li><li>3. Huang, H.Y. <i>et al.</i> (2008) Immunohistochemical and biogenetic features of diffuse-type tenosynovial giant cell tumors: the potential roles of cyclin A, P53, and deletion of 15q in sarcomatous transformation. <a href="#">Clin Cancer Res. 14: 6023-32.</a></li><li>4. Iannone, F. <i>et al.</i> (2005) Increased Bcl-2/p53 ratio in human osteoarthritic cartilage: a possible role in regulation of chondrocyte metabolism. <a href="#">Ann Rheum Dis. 64: 217-21.</a></li><li>5. Lin, L.C. <i>et al.</i> (2006) p53 and p27 as predictors of clinical outcome for rectal-cancer patients receiving neoadjuvant therapy. <a href="#">Surg Oncol. 15: 211-6.</a></li></ol>
<b>Storage</b>	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1703">https://www.bio-rad-antibodies.com/SDS/MCA1703</a> 10040
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

## Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL \(MCA691\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M365602:200529'

**Printed on 29 Feb 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)