

## Datasheet: MCA1710AMO

**BATCH NUMBER 149462**

<b>Description:</b>	MOUSE ANTI HUMAN CD20:Amethyst Orange
<b>Specificity:</b>	CD20
<b>Format:</b>	Amethyst Orange
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	2H7
<b>Isotype:</b>	IgG2b
<b>Quantity:</b>	0.1 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

#### Target Species

Human

#### Species Cross Reactivity

Reacts with: Rhesus Monkey

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

#### Product Form

Purified IgG conjugated to Amethyst Orange - liquid

#### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
Amethyst Orange	405	540

#### Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

#### Buffer Solution

Phosphate buffered saline

<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P11836</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">931</a>    MS4A1    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	CD20
<b>Specificity</b>	<p><b>Mouse anti Human CD20 antibody, clone 2H7</b> recognizes the human CD20 cell surface antigen, a 33-37 kDa non-glycosylated phosphoprotein.</p> <p>The CD20 antigen is expressed during pre-B-cell development. It is present on both resting and activated B-cells but is lost prior to terminal B-cell differentiation into plasma cells.</p> <p>The epitope recognized by clone 2H7 has been mapped to the following sequence found in the large extracellular loop of human CD20: YNCEPANPSEKNSPST. Furthermore it appears that Mouse anti Human CD20 antibody, clone 2H7 only recognizes human CD20 in its native oligomeric form (<a href="#">Polyak et al. 2002</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells or 100ul whole blood
<b>References</b>	<ol style="list-style-type: none"> <li>Chan, H.T. <i>et al.</i> (2003) CD20-induced lymphoma cell death is independent of both caspases and its redistribution into triton X-100 insoluble membrane rafts. <a href="#">Cancer Res. 63: 5480-9.</a></li> <li>Cragg, M.S. <i>et al.</i> (2003) Complement-mediated lysis by anti-CD20 mAb correlates with segregation into lipid rafts. <a href="#">Blood. 101: 1045-52.</a></li> <li>Jaramillo, M.C. <i>et al.</i> (2009) Increased manganese superoxide dismutase expression or treatment with manganese porphyrin potentiates dexamethasone-induced apoptosis in lymphoma cells. <a href="#">Cancer Res. 69: 5450-7.</a></li> <li>Teeling, J.L. <i>et al.</i> (2006) The biological activity of human CD20 monoclonal antibodies is linked to unique epitopes on CD20. <a href="#">J Immunol. 177 (1): 362-71.</a></li> <li>Polyak, M.J. <i>et al.</i> (2002) Alanine-170 and proline-172 are critical determinants for extracellular CD20 epitopes; heterogeneity in the fine specificity of CD20 monoclonal antibodies is defined by additional requirements imposed by both amino acid sequence and quaternary structure. <a href="#">Blood. 1;99:3256-62.</a></li> <li>Greig, B. <i>et al.</i> (2014) Stabilization media increases recovery in paucicellular cerebrospinal fluid specimens submitted for flow cytometry testing. <a href="#">Cytometry B Clin Cytom. 86: 135-8.</a></li> <li>van den Akker, E. <i>et al.</i> (2010) The majority of the in vitro erythroid expansion potential resides in CD34(-) cells, outweighing the contribution of CD34(+) cells and significantly</li> </ol>

increasing the erythroblast yield from peripheral blood samples. [Haematologica. 95: 1594-8.](#)

8. Jaramillo, M.C. *et al.* (2015) Manganese (III) meso-tetrakis N-ethylpyridinium-2-yl porphyrin acts as a pro-oxidant to inhibit electron transport chain proteins, modulate bioenergetics, and enhance the response to chemotherapy in lymphoma cells. [Free Radic Biol Med. 83: 89-100.](#)

9. Cecchinato, V. *et al.* (2017) Impairment of CCR6+ and CXCR3+ Th Cell Migration in HIV-1 Infection Is Rescued by Modulating Actin Polymerization. [J Immunol. 198 \(1\): 184-195.](#)

10. Kohler, S.L. *et al.* (2016) Germinal Center T Follicular Helper Cells Are Highly Permissive to HIV-1 and Alter Their Phenotype during Virus Replication. [J Immunol. 196 \(6\): 2711-22.](#)

11. Grobárová V *et al.* (2016) Quambalarine B, a Secondary Metabolite from *Quambalaria cyanescens* with Potential Anticancer Properties. [J Nat Prod. 79 \(9\): 2304-14.](#)

12. Popov, J. *et al.* (2017) Unique therapeutic properties and preparation methodology of multivalent rituximab-lipid nanoparticles. [Eur J Pharm Biopharm. 117: 256-69.](#)

13. Sieg, M. *et al.* (2019) A New Genotype of Feline Morbillivirus Infects Primary Cells of the Lung, Kidney, Brain and Peripheral Blood. [Viruses. 11 \(2\) Feb 09 \[Epub ahead of print\].](#)

---

<b>Storage</b>	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
----------------	---

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1710AMO">https://www.bio-rad-antibodies.com/SDS/MCA1710AMO</a> 10041
--------------------------------------	---

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

## Related Products

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M360646:191223'

Printed on 10 May 2024

