

Datasheet: MCA1710SBV515

BATCH NUMBER 100004361

Description:	MOUSE ANTI HUMAN CD20:StarBright Violet 515
Specificity:	CD20
Format:	StarBright Violet 515
Product Type:	Monoclonal Antibody
Clone:	2H7
Isotype:	IgG2b
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species Human

Species Cross Reactivity Reacts with: Rhesus Monkey
N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form Purified IgG conjugated to StarBright Violet 515 - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	StarBright Violet 515	401	516

Preparation Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative	0.09% Sodium Azide (NaN ₃)
Stabilisers	1% Bovine Serum Albumin 0.1% Pluronic F68 0.1% PEG 3350

External Database Links

UniProt:

[P11836](#) [Related reagents](#)

Entrez Gene:

[931](#) MS4A1 [Related reagents](#)

Synonyms CD20

Specificity **Mouse anti Human CD20 antibody, clone 2H7** recognizes the human CD20 cell surface antigen, a 33-37 kDa non-glycosylated phosphoprotein.

The CD20 antigen is expressed during pre-B-cell development. It is present on both resting and activated B-cells but is lost prior to terminal B-cell differentiation into plasma cells.

The epitope recognized by clone 2H7 has been mapped to the following sequence found in the large extracellular loop of human CD20: YNCEPANPSEKNPST. Furthermore it appears that Mouse anti Human CD20 antibody, clone 2H7 only recognizes human CD20 in its native oligomeric form ([Polyak *et al.* 2002](#)).

Flow Cytometry Use 5ul of the suggested working dilution to label 10⁶ cells in 100ul. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

- References**
1. Chan, H.T. *et al.* (2003) CD20-induced lymphoma cell death is independent of both caspases and its redistribution into triton X-100 insoluble membrane rafts. [Cancer Res. 63: 5480-9.](#)
 2. Cragg, M.S. *et al.* (2003) Complement-mediated lysis by anti-CD20 mAb correlates with segregation into lipid rafts. [Blood. 101: 1045-52.](#)
 3. Jaramillo, M.C. *et al.* (2009) Increased manganese superoxide dismutase expression or treatment with manganese porphyrin potentiates dexamethasone-induced apoptosis in lymphoma cells. [Cancer Res. 69: 5450-7.](#)
 4. Teeling, J.L. *et al.* (2006) The biological activity of human CD20 monoclonal antibodies is linked to unique epitopes on CD20. [J Immunol. 177 \(1\): 362-71.](#)
 5. Polyak, M.J. *et al.* (2002) Alanine-170 and proline-172 are critical determinants for extracellular CD20 epitopes; heterogeneity in the fine specificity of CD20 monoclonal antibodies is defined by additional requirements imposed by both amino acid sequence and quaternary structure. [Blood. 1;99:3256-62.](#)
 6. Greig, B. *et al.* (2014) Stabilization media increases recovery in paucicellular cerebrospinal fluid specimens submitted for flow cytometry testing. [Cytometry B Clin Cytom. 86: 135-8.](#)
 7. van den Akker, E. *et al.* (2010) The majority of the in vitro erythroid expansion potential resides in CD34(-) cells, outweighing the contribution of CD34(+) cells and significantly

increasing the erythroblast yield from peripheral blood samples. [Haematologica. 95: 1594-8.](#)

8. Jaramillo, M.C. *et al.* (2015) Manganese (III) meso-tetrakis N-ethylpyridinium-2-yl porphyrin acts as a pro-oxidant to inhibit electron transport chain proteins, modulate bioenergetics, and enhance the response to chemotherapy in lymphoma cells. [Free Radic Biol Med. 83: 89-100.](#)

9. Cecchinato, V. *et al.* (2017) Impairment of CCR6+ and CXCR3+ Th Cell Migration in HIV-1 Infection Is Rescued by Modulating Actin Polymerization. [J Immunol. 198 \(1\): 184-195.](#)

10. Kohler, S.L. *et al.* (2016) Germinal Center T Follicular Helper Cells Are Highly Permissive to HIV-1 and Alter Their Phenotype during Virus Replication. [J Immunol. 196 \(6\): 2711-22.](#)

11. Grobárová V *et al.* (2016) Quambalarine B, a Secondary Metabolite from *Quambalaria cyanescens* with Potential Anticancer Properties. [J Nat Prod. 79 \(9\): 2304-14.](#)

12. Popov, J. *et al.* (2017) Unique therapeutic properties and preparation methodology of multivalent rituximab-lipid nanoparticles. [Eur J Pharm Biopharm. 117: 256-69.](#)

13. Sieg, M. *et al.* (2019) A New Genotype of Feline Morbillivirus Infects Primary Cells of the Lung, Kidney, Brain and Peripheral Blood. [Viruses. 11 \(2\) Feb 09 \[Epub ahead of print\].](#)

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
----------------	---

Guarantee	12 months from date of despatch
------------------	---------------------------------

Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
-------------------------	---

Health And Safety Information	Material Safety Datasheet documentation #20438 available at: https://www.bio-rad-antibodies.com/SDS/MCA1710SBV515 20438
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M373242:200901'

Printed on 25 Mar 2023