

Datasheet: MCA1719F BATCH NUMBER 166797

Description:	MOUSE ANTI HUMAN CD45:FITC
Specificity:	CD45
Other names:	LCA
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	HI30
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			1/5

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human			
Species Cross	Reacts with: Chim	panzee		
Reactivity	N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated fo further information.			viewed publications or
Product Form	Purified IgG conju	gated to Fluorescein Isotl	niocyanate Isomer 1	(FITC) - liquid
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	FITO	400	505	

	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography		aphy
Buffer Solution	Phosphate buffered sa	aline	

Preservative Stabilisers	0.09% sodium azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml
External Database Links	UniProt: P08575 Related reagents
	Entrez Gene: <u>5788</u> PTPRC <u>Related reagents</u>
Synonyms	CD45
Specificity	Mouse anti Human CD45 antibody, clone HI30 recognizes the human CD45 cell surface antigen, also known as the leukocyte common antigen.
	CD45 is expressed by all leukocytes; expression on granulocytes being weaker than on lymphocytes.
Flow Cytometry	Use 10μl of the suggested working dilution to label 10 ⁶ cells or 100μl whole blood
References	1. Eksteen, B. <i>et al.</i> (2009) Gut homing receptors on CD8 T cells are retinoic acid dependent and not maintained by liver dendritic or stellate cells. <u>Gastroenterology. 137</u> (1): 320-9.
	 Alvarez-gonzalez, C. et al. (2013) Cord blood Lin(-)CD45(-) embryonic-like stem cells are a heterogeneous population that lack self-renewal capacity. PLoS One. 8 (6): e67968. Tripathy, N.K. et al. (2018) Cardiomyogenic Heterogeneity of Clonal Subpopulations of Human Bone Marrow Mesenchymal Stem Cells. J Stem Cells Regen Med. 14 (1): 27-33.
Storage	Store at +4°C. DO NOT FREEZE.
	This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1719F 10040
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M411192:221102'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint