# Datasheet: MCA1723 BATCH NUMBER 163758

Description:	MOUSE ANTI HUMAN CD8 BETA
Specificity:	CD8 BETA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	2ST8.5H7
Isotype:	lgG2a
Quantity:	0.2 mg

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is						
	derived from testing within our laboratories, peer-reviewed publications or personal						
	communications from the originators. Please refer to references indicated for further						
	information. For general	protocol re	ecommen	dations, please visit <u>w</u>	ww.bio-		
	rad-antibodies.com/protocols.						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	-			Neat - 1/10		
	Immunohistology - Frozen			•			
	Immunohistology - Paraffin			•			
	ELISA			•			
	Immunoprecipitation	-					
	Western Blotting	•					
	Where this antibody has not been tested for use in a particular technique this does not						
	necessarily exclude its use in such procedures. Suggested working dilutions are given as						
	a quide only. It is recomn	nended th	at the use	er titrates the antibody	for use in their own		
	system using appropriate	e negative	/positive o	controls.			
Target Species	Human						
Species Cross	Reacts with: Cynomolgue	s monkey,	Rhesus I	Monkey			
Reactivity	N.B. Antibody reactivity and working conditions may vary between species. Cross						
	reactivity is derived from	testing wi	thin our la	aboratories, peer-reviev	wed publications or		
	personal communications	s from the	originato	rs. Please refer to refe	rences indicated for		
	further information.		-				
Product Form	Purified IgG - lyophilized						
Reconstitution	Reconstitute with 0.2ml c	listilled wa	ater				
	Care should be taken during reconstitution as the protein may appear as a film at the						

	bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution. For long term storage the addition of 0.09% sodium azide is recommended. N.B. For functional studies do not add sodium azide
Preparation	Purified IgG prepared by affinity chromatography from Ascites
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Normal human T lymphocytes.
External Database Links	UniProt: <u>P10966</u> <u>Related reagents</u>
	<u>926</u> CD8B <u>Related reagents</u>
Synonyms	CD8B1
RRID	AB_322685
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
Specificity	<b>Mouse anti Human CD8 beta antibody, clone 2ST8.5H7</b> recognizes the human CD8 cell surface antigen. The antibody binds to the beta chain, recognizing an epitope that is dependent upon expression of both CD8 alpha and CD8 beta.
	CD8 molecules are composed of dimers of either alpha/alpha or alpha/beta polypeptide combinations. Both types of dimer may co-exist at the cell surface. Clone 2ST8.5H7 is also referred to as clone T8/2T8-5H7 and binding is dependent on the expression of the alpha/beta dimer.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
References	<ol> <li>Terry, L.A. <i>et al.</i> (1990) Differential expression and regulation of the human CD8 alpha and CD8 beta chains. <u>Tissue Antigens 35: 82-91.</u></li> <li>DiSanto, J.P. <i>et al.</i> (1991) Generation of anti-human CD8 beta-specific antibodies using transfectants expressing mixed-species CD8 heterodimers. <u>J Immunol Methods. 141 (1):</u> <u>123-31.</u></li> <li>de la Calle-Martin, O. <i>et al.</i> (2001) Familial CD8 deficiency due to a mutation in the CD8 alpha gene. <u>J Clin Invest. 108: 117-23.</u></li> <li>Schmitz, J.E. <i>et al</i> (1998) Expression of the CD8alpha beta-heterodimer on CD8(+) T</li> </ol>

	lymphocytes in peripheral blood lymphocytes of human immunodeficiency virus- and human immunodeficiency virus+ individuals. <u>Blood. 92: 198-206.</u>			
Storage	Prior to reconstitution store at +4°C. After reconstitution store at -20°C. Storage in frost-free freezers is not recommended. This product should be stored			
	this product contain a precipitate we recommend microcentrifugation before use.	IU		
Guarantee	Guaranteed until date of expiry. Please see product label.			
Health And Safety Information	Material Safety Datasheet documentation #10294 available at: https://www.bio-rad-antibodies.com/SDS/MCA1723 10294			
Regulatory	For research purposes only			

### **Related Products**

### **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12)	RPE	
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>		
Goat Anti Mouse IgG (STAR76)	RPE	
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>	
Rabbit Anti Mouse IgG (STAR13)	HRP	
Goat Anti Mouse IgG (Fc) (STAR120)	<u>FITC, HRP</u>	
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>	
Goat Anti Mouse IgG (STAR77)	HRP	
Goat Anti Mouse IgG (H/L) (STAR117)	<u>Alk. Phos.</u> , <u>DyLight®488</u> , <u>DyLight®550</u> ,	
	DyLight®650, DyLight®680, DyLight®800,	
	<u>FITC, HRP</u>	

#### **Recommended Negative Controls**

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad	.com	Email: antibody_sales_uk@bio-rad	.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M390947:211008'

#### Printed on 10 Mar 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint