

Datasheet: MCA1724PET

Description:	MOUSE ANTI HUMAN CD152:RPE
Specificity:	CD152
Other names:	CTLA-4
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	BNI3
Isotype:	IgG2a
Quantity:	25 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) **Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.**

Target Species	Human		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein A from ascites		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	0.2% Bovine Serum Albumin		
Immunogen	Human CTLA-4/human IgG heavy chain fusion protein.		
External Database Links	UniProt: P16410 Related reagents		
	Entrez Gene:		

Synonyms	CD152
-----------------	-------

Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse P3/X63-Ag8.653 myeloma cell line.
------------------------	--

Specificity	<p>Mouse anti Human CD152 antibody, clone BNI3 recognizes human CD152, also known as CTLA-4 (cytotoxic T-lymphocyte-associated antigen 4), an inhibitory receptor and negative regulator of T-cell responses. CD152 is a single pass type 1 transmembrane protein belonging to the immunoglobulin superfamily containing a single Ig-v-like domain in the extracellular region.</p> <p>CD152 along with CD28 binds to the co-stimulatory molecules CD80 and CD86 (Azuma <i>et al.</i> 1993).</p> <p>Mouse anti human CD152 antibody, clone BNI3 is able to block ligand binding on the Raji B-cell line (Steiner <i>et al.</i> 2001) and blocks binding of an alternative clone, BNI8 to CTLA-4/Ig in ELISA. Mouse anti Human CD152 antibody, clone BNI3 binds to the same epitope as classified anti CTLA-4 clones 11D4 and 10A8 (Wang <i>et al.</i> In: Leukocyte typing VI 1997 Garland Publishing Inc. pp97-98, Bull World Health Organ. 1997).</p> <p>The cytoplasmic domain of CD152 contains a critical tyrosine at residue 201 phosphorylated by Janus Kinase 2 which subsequently controls surface expression through regulation of CD152 interaction with AP-2 (Shiratori <i>et al.</i> 1997, Chikuma <i>et al.</i> 2000). CD152 is expressed primarily as an intracellular antigen with transport to the cell surface under tight regulation of several molecules including Trim, PLD and TIRC7, CD152 also demonstrates rapid internalization once expressed at the cell surface (Valk <i>et al.</i> 2008). CD152 plays a significant role in maintaining tolerance to self antigens and defects in CD152 presentation and expression has been implicated in a number of autoimmune diseases (Romo-Tena <i>et al.</i> 2013).</p>
--------------------	---

Flow Cytometry	Use 20ul of the suggested working dilution to label 5x10 ⁵ cells in 100ul.
-----------------------	---

References	<ol style="list-style-type: none">1. Linsley, P.S. <i>et al.</i> (1992) Coexpression and functional cooperation of CTLA-4 and CD28 on activated T lymphocytes. J Exp Med. 176 (6): 1595-604.2. Kuiper, H.M. <i>et al.</i> (1995) Activated T cells can induce high levels of CTLA-4 expression on B cells. J Immunol. 155 (4): 1776-83.3. Castan, J. <i>et al.</i> (1997) Accumulation of CTLA-4 expressing T lymphocytes in the germinal centres of human lymphoid tissues. Immunology. 90 (2): 265-71.4. Lee, C.C. <i>et al.</i> (2009) The regulatory function of umbilical cord blood CD4(+) CD25(+) T cells stimulated with anti-CD3/anti-CD28 and exogenous interleukin (IL)-2 or IL-15. Pediatr Allergy Immunol. 20 (7): 624-32.5. Pistillo, M.P. <i>et al.</i> (2003) CTLA-4 is not restricted to the lymphoid cell lineage and can function as a target molecule for apoptosis induction of leukemic cells. Blood. 101: 202-9.6. Tan, P.H. <i>et al.</i> (2005) Creation of tolerogenic human dendritic cells via intracellular CTLA4: a novel strategy with potential in clinical immunosuppression. Blood. 106: 2936-43.7. Steiner, K. <i>et al.</i> (2001) Increased expression of CTLA-4 (CD152) by T and B lymphocytes in Wegener's granulomatosis. Clin Exp Immunol. 126: 143-50.8. Lu, C.H. <i>et al.</i> (2016) DNA Methyltransferase Inhibitor Promotes Human CD4⁺CD25^hFOXP3⁺ Regulatory T Lymphocyte Induction under Suboptimal TCR Stimulation. Front Immunol. 7: 488.9. Steiner, K. <i>et al.</i> (1999) Enhanced expression of CTLA-4 (CD152) on CD4⁺ T cells in HIV infection. Clin Exp Immunol. 115 (3): 451-7.10. Ward, F.J. <i>et al.</i> (2013) The soluble isoform of CTLA-4 as a regulator of T-cell responses. Eur J
-------------------	---

Further Reading	1. Chin, L.T. <i>et al.</i> (2008) Immune intervention with monoclonal antibodies targeting CD152 (CTLA-4) for autoimmune and malignant diseases. Chang Gung Med J. 31: 1-15.
------------------------	---

Storage	Store at +4°C. DO NOT FREEZE This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
----------------	---

Shelf Life	18 months from date of despatch.
-------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:RPE \(MCA929PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M322968:180727'

Printed on 06 Dec 2018