

Datasheet: MCA1738

BATCH NUMBER 166938

Description:	MOUSE ANTI HUMAN CD31
Specificity:	CD31
Other names:	PECAM-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	WM59
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Cynomolgus monkey, Rhesus Monkey</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
External Database Links	<p>UniProt: P16284 Related reagents</p> <p>Entrez Gene: 5175 PECAM1 Related reagents</p>
RRID	AB_322710
Specificity	<p>Mouse anti Human CD31 monoclonal antibody, clone WM59 recognizes the human CD31 antigen, a ~130 kDa single pass type I transmembrane glycoprotein bearing six C2 immunoglobulin domains. CD31 is expressed by all continuous endothelia including arteries, veins and non-sinusoidal capillaries, platelets, granulocytes and some lymphocytes. CD31 is not expressed by discontinuous endothelia such as hepatic sinusoids and splenic red pulp (Muller et al. 1989). CD31 is also known as PECAM-1.</p> <p>The binding epitope for mouse anti human CD31, clone WM59 has been mapped to the Ig-like domain 2 (Fawcett et al. 1995).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> Paul, G. <i>et al.</i> (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. PLoS One. 7: e35577. Urquhart, P. <i>et al.</i> (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. J Pharmacol Exp Ther. 321 (2): 656-62. Reedquist, K.A. <i>et al.</i> (2000) The small GTPase, Rap1, mediates CD31-induced integrin adhesion. J Cell Biol. 148: 1151-8. Vernon-Wilson, E.F. <i>et al.</i> (2007) CD31 delays phagocyte membrane repolarization to promote efficient binding of apoptotic cells. J Leukoc Biol. 82: 1278-88. Hilbe W <i>et al.</i> (2003) Immunohistochemical typing of non-small cell lung cancer on cryostat sections: correlation with clinical parameters and prognosis. J Clin Pathol. 56 (10): 736-41. Stein, A. <i>et al.</i> (2010) Local erythropoietin and endothelial progenitor cells improve regional cardiac function in acute myocardial infarction. BMC Cardiovasc Disord. Sep; 10:43. Woollard, K.J. <i>et al.</i> (2002) Direct modulatory effect of C-reactive protein on primary

- human monocyte adhesion to human endothelial cells. [Clin Exp Immunol. 130: 256-62.](#)
8. Theberge, A.B. *et al.* (2015) Microfluidic multiculture assay to analyze biomolecular signaling in angiogenesis. [Anal Chem. 87 \(6\): 3239-46.](#)
9. Hilbe W *et al.* (2004) CD133 positive endothelial progenitor cells contribute to the tumour vasculature in non-small cell lung cancer. [J Clin Pathol. 57 \(9\): 965-9.](#)
10. Palakkan, A.A. *et al.* (2015) Polarisation and functional characterisation of hepatocytes derived from human embryonic and mesenchymal stem cells. [Biomed Rep. 3 \(5\): 626-636.](#)
11. Newey SE *et al.* (2014) The hematopoietic chemokine CXCL12 promotes integration of human endothelial colony forming cell-derived cells into immature vessel networks. [Stem Cells Dev. 23 \(22\): 2730-43.](#)
12. Fabre-Mersseman V *et al.* (2011) CD4⁺ recent thymic emigrants are infected by HIV in vivo, implication for pathogenesis. [AIDS. 25 \(9\): 1153-62.](#)
13. Patten PE *et al.* (2008) CD38 expression in chronic lymphocytic leukemia is regulated by the tumor microenvironment. [Blood. 111 \(10\): 5173-81.](#)
14. Katz SC *et al.* (2004) Liver sinusoidal endothelial cells are insufficient to activate T cells. [J Immunol. 173 \(1\): 230-5.](#)
15. Pfisterer K *et al.* (2015) CD90(+) human dermal stromal cells are potent inducers of FoxP3(+) regulatory T cells. [J Invest Dermatol. 135 \(1\): 130-41.](#)
16. Hale, S.J. *et al.* (2015) CXCR2 modulates bone marrow vascular repair and haematopoietic recovery post-transplant. [Br J Haematol. 169 \(4\): 552-64.](#)
17. Muthana, M. *et al.* (2015) Directing cell therapy to anatomic target sites in vivo with magnetic resonance targeting. [Nat Commun. 6: 8009.](#)
18. Schuster, C. *et al.* (2015) Development of Blood and Lymphatic Endothelial Cells in Embryonic and Fetal Human Skin. [Am J Pathol. 185 \(9\): 2563-74.](#)
19. Somers, E. *et al.* (2016) Vascular Defects and Spinal Cord Hypoxia in Spinal Muscular Atrophy. [Ann Neurol. 79 \(2\): 217-30.](#)
20. Soh, B.S. *et al.* (2016) Endothelin-1 supports clonal derivation and expansion of cardiovascular progenitors derived from human embryonic stem cells. [Nat Commun. 7: 10774.](#)
21. GarikipatiV, N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. [PLoS One. 13 \(2\): e0192244.](#)
22. Duque, J.C. *et al.* (2019) Vascularization of the arteriovenous fistula wall and association with maturation outcomes. [J Vasc Access. : 1129729819863584. \[Epub ahead of print\]](#)
23. Kim, J.S. *et al.* (2021) Randomization to Omega-3 Fatty Acid Supplementation and Endothelial Function in COPD: The COD-Fish Randomized Controlled Trial. [Chronic Obstr Pulm Dis. 8\(1\): 41-53.](#)
24. Bye, A.P. *et al.* (2018) Immobilization of Nonactivated Unfixed Platelets for Real-Time Single-Cell Analysis. [Methods Mol Biol. 1812: 1-11.](#)
25. Chai, S. *et al.* (2022) Identification of epithelial and mesenchymal circulating tumor cells in clonal lineage of an aggressive prostate cancer case. [NPJ Precis Oncol. 6 \(1\): 41.](#)
26. Bettin, L. *et al.* (2023) Co-stimulation by TLR7/8 ligand R848 modulates IFN- γ production of porcine $\gamma\delta$ T cells in a microenvironment-dependent manner. [Dev Comp Immunol. 138: 104543.](#)
27. Seo, J. *et al.* (2023) Plasticity of circulating tumor cells in small cell lung cancer. [Sci Rep. 13 \(1\): 11775.](#)

28. Shishido, S.N. *et al.* (2024) Cancer-related cells and oncosomes in the liquid biopsy of pancreatic cancer patients undergoing surgery. [NPJ Precis Oncol. 8 \(1\): 36.](#)

Further Reading	1. DeLisser, H.M. <i>et al.</i> (1994) Molecular and functional aspects of PECAM-1/CD31. Immunol Today. 15 (10): 490-5.
Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1738 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M404544:221102'

Printed on 08 Mar 2024