

Datasheet: MCA1738B

Description:	MOUSE ANTI HUMAN CD31:Biotin
Specificity:	CD31
Other names:	PECAM-1
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	WM59
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

RRID AB_322533

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Human

Species Cross Reactivity Reacts with: Cynomolgus monkey, Rhesus Monkey
N.B. Antibody reactivity and working conditions may vary between species.

Product Form Purified IgG conjugated to Biotin - liquid

Preparation Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide
Stabilisers 1% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.1mg/ml

External Database Links
UniProt:
[P16284](#) [Related reagents](#)

Entrez Gene:

[5175](#) PECAM1 [Related reagents](#)

Specificity

Mouse anti Human CD31 monoclonal antibody, clone WM59 recognizes the human CD31 antigen, a ~130 kDa single pass type I transmembrane glycoprotein bearing six [C2 immunoglobulin domains](#). CD31 is expressed by all continuous endothelia including arteries, veins and non-sinusoidal capillaries, platelets, granulocytes and some lymphocytes. CD31 is not expressed by discontinuous endothelia such as hepatic sinusoids and splenic red pulp ([Muller et al. 1989](#)). CD31 is also known as PECAM-1.

The binding epitope for mouse anti human CD31, clone WM59 has been mapped to the Ig-like domain 2 ([Fawcett et al. 1995](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10^6 cells or 100ul whole blood

References

1. Cabañas, C. *et al.* (1989) Characterization of a novel myeloid antigen regulated during differentiation of monocytic cells. [Eur J Immunol. 19 \(8\): 1373-8.](#)
2. Paul, G. *et al.* (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. [PLoS One. 7: e35577.](#)
3. Stockinger, H. *et al.* (1990) Molecular characterization and functional analysis of the leukocyte surface protein CD31. [J Immunol. 145 \(11\): 3889-97.](#)
4. DeLisser, H.M. *et al.* (1994) Molecular and functional aspects of PECAM-1/CD31. [Immunol Today. 15 \(10\): 490-5.](#)
5. Urquhart, P. *et al.* (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. [J Pharmacol Exp Ther. 321 \(2\): 656-62.](#)
6. Fawcett, J. *et al.* (1995) Mapping the homotypic binding sites in CD31 and the role of CD31 adhesion in the formation of interendothelial cell contacts. [J Cell Biol. 128:1229-41.](#)
7. Vernon-Wilson, E.F. *et al.* (2007) CD31 delays phagocyte membrane repolarization to promote efficient binding of apoptotic cells. [J Leukoc Biol. 82: 1278-88.](#)
8. Johnston, A. *et al.* (2005) The anti-inflammatory action of methotrexate is not mediated by lymphocyte apoptosis, but by the suppression of activation and adhesion molecules. [Clin Immunol. 114: 154-63.](#)
9. Reedquist, K.A. *et al.* (2000) The small GTPase, Rap1, mediates CD31-induced integrin adhesion. [J Cell Biol. 148: 1151-8.](#)
10. Stein, A. *et al.* (2010) Local erythropoietin and endothelial progenitor cells improve regional cardiac function in acute myocardial infarction. [BMC Cardiovasc Disord. Sep; 10:43.](#)
11. Woollard, K.J. *et al.* (2002) Direct modulatory effect of C-reactive protein on primary human monocyte adhesion to human endothelial cells. [Clin Exp Immunol. 130: 256-62.](#)
12. Theberge, A.B. *et al.* (2015) Microfluidic multiculture assay to analyze biomolecular signaling in angiogenesis. [Anal Chem. 87 \(6\): 3239-46.](#)
13. Hilbe W *et al.* (2004) CD133 positive endothelial progenitor cells contribute to the tumour vasculature in non-small cell lung cancer. [J Clin Pathol. 57 \(9\): 965-9.](#)
14. Hilbe W *et al.* (2003) Immunohistochemical typing of non-small cell lung cancer on cryostat sections: correlation with clinical parameters and prognosis. [J Clin Pathol. 56 \(10\): 736-41.](#)
15. Palakkan, A.A. *et al.* (2015) Polarisation and functional characterisation of hepatocytes derived from human embryonic and mesenchymal stem cells. [Biomed Rep. 3 \(5\): 626-636.](#)
16. Newey SE *et al.* (2014) The hematopoietic chemokine CXCL12 promotes integration of human endothelial colony forming cell-derived cells into immature vessel networks. [Stem Cells Dev. 23 \(22\): 2730-43.](#)
17. Fabre-Mersseman V *et al.* (2011) CD4⁺ recent thymic emigrants are infected by HIV in vivo, implication for pathogenesis. [AIDS. 25 \(9\): 1153-62.](#)
18. Patten PE *et al.* (2008) CD38 expression in chronic lymphocytic leukemia is regulated by the

- tumor microenvironment. [Blood. 111 \(10\): 5173-81.](#)
19. Katz SC *et al.* (2004) Liver sinusoidal endothelial cells are insufficient to activate T cells. [J Immunol. 173 \(1\): 230-5.](#)
20. Pfisterer K *et al.* (2015) CD90(+) human dermal stromal cells are potent inducers of FoxP3(+) regulatory T cells. [J Invest Dermatol. 135 \(1\): 130-41.](#)
21. Hale, S.J. *et al.* (2015) CXCR2 modulates bone marrow vascular repair and haematopoietic recovery post-transplant. [Br J Haematol. 169 \(4\): 552-64.](#)
22. Muthana, M. *et al.* (2015) Directing cell therapy to anatomic target sites in vivo with magnetic resonance targeting. [Nat Commun. 6: 8009.](#)
23. Schuster, C. *et al.* (2015) Development of Blood and Lymphatic Endothelial Cells in Embryonic and Fetal Human Skin. [Am J Pathol. 185 \(9\): 2563-74.](#)
24. Somers, E. *et al.* (2016) Vascular Defects and Spinal Cord Hypoxia in Spinal Muscular Atrophy. [Ann Neurol. 79 \(2\): 217-30.](#)
25. Soh, B.S. *et al.* (2016) Endothelin-1 supports clonal derivation and expansion of cardiovascular progenitors derived from human embryonic stem cells. [Nat Commun. 7: 10774.](#)
26. Yi, T. *et al.* (2015) Manufacture of Clinical-Grade Human Clonal Mesenchymal Stem Cell Products from Single Colony Forming Unit-Derived Colonies Based on the Subfractionation Culturing Method. [Tissue Eng Part C Methods. 21 \(12\): 1251-62.](#)
27. GarikipatiV, N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. [PLoS One. 13 \(2\): e0192244.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory

For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South

Tel: +1 800 265 7376

America

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M322981:180727'

Printed on 10 Dec 2018