

Datasheet: MCA1738F BATCH NUMBER 154447

| Description:  | MOUSE ANTI HUMAN CD31:FITC |
|---------------|----------------------------|
| Specificity:  | CD31                       |
| Other names:  | PECAM-1                    |
| Format:       | FITC                       |
| Product Type: | Monoclonal Antibody        |
| Clone:        | WM59                       |
| Isotype:      | IgG1                       |
| Quantity:     | 0.1 mg                     |
|               |                            |

### **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | -   |    |                | Neat - 1/2         |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| <b>Species Cross</b> |
|----------------------|
| Reactivity           |

**Target Species** 

Human

Reacts with: Cynomolgus monkey, Rhesus Monkey

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

| _ |    | -   |     | _                         |        |
|---|----|-----|-----|---------------------------|--------|
| D | ra | d., | nt. | $\mathbf{E}_{\mathbf{A}}$ | rm     |
| г | ľ  | uu  | Lι  | гυ                        | ,, ,,, |

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) |  |
|-----------|-------------|---------------------|-------------------|--|
|           | FITC        | 490                 | 525               |  |
|           |             |                     |                   |  |

### Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

| Buffer Solution                   | Phosphate buffered saline   |
|-----------------------------------|---|
| Preservative<br>Stabilisers       | 0.09% Sodium Azide 1% Bovine Serum Albumin  |
| Approx. Protein<br>Concentrations | IgG concentration 0.1mg/ml  |
| External Database<br>Links        | UniProt: P16284 Related reagents  Entrez Gene: 5175 PECAM1 Related reagents   |
| RRID                              | AB_322848   |
| Specificity                       | Mouse anti Human CD31 monoclonal antibody, clone WM59 recognizes the human CD31 antigen, a ~130 kDa single pass type I transmembrane glycoprotein bearing six C2 immunoglobulin domains. CD31 is expressed by all continuous endothelia including arteries, veins and non-sinusoidal capillaries, platelets,granulocytes and some lymphocytes. CD31 is not expressed by discontinuous endothelia such as hepatic sinusoids and splenic red pulp (Muller et al. 1989).CD31 is also known as PECAM-1. |

The binding epitope for mouse anti human CD31, clone WM59 has been mapped to the Ig-like domain 2 (Fawcett et al. 1995).

#### Flow Cytometry

Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells or 100ul whole blood

#### References

- 1. Paul, G. et al. (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. PLoS One. 7: e35577.
- 2. Stockinger, H. et al. (1990) Molecular characterization and functional analysis of the leukocyte surface protein CD31. J Immunol. 145 (11): 3889-97.
- 3. DeLisser, H.M. et al. (1994) Molecular and functional aspects of PECAM-1/CD31. Immunol Today. 15 (10): 490-5.
- 4. Urguhart, P. et al. (2007) Carbon monoxide-releasing molecules modulate leukocyteendothelial interactions under flow. J Pharmacol Exp Ther. 321 (2): 656-62.
- 5. Reedquist, K.A. et al. (2000) The small GTPase, Rap1, mediates CD31-induced integrin adhesion. J Cell Biol. 148: 1151-8.
- 6. Vernon-Wilson, E.F. et al. (2007) CD31 delays phagocyte membrane repolarization to promote efficient binding of apoptotic cells. <u>J Leukoc Biol. 82: 1278-88.</u>
- 7. Johnston, A. et al. (2005) The anti-inflammatory action of methotrexate is not mediated by lymphocyte apoptosis, but by the suppression of activation and adhesion molecules. Clin Immunol. 114: 154-63.
- 8. Hilbe W et al. (2003) Immunohistochemical typing of non-small cell lung cancer on cryostat sections: correlation with clinical parameters and prognosis. J Clin Pathol. 56 (10): 736-41.
- 9. Stein, A. et al. (2010) Local erythropoietin and endothelial progenitor cells improve

regional cardiac function in acute myocardial infarction. <u>BMC Cardiovasc Disord. Sep;</u> 10:43.

- 10. Woollard, K.J. *et al.* (2002) Direct modulatory effect of C-reactive protein on primary human monocyte adhesion to human endothelial cells. <u>Clin Exp Immunol</u>. 130: 256-62.
- 11. Theberge, A.B. *et al.* (2015) Microfluidic multiculture assay to analyze biomolecular signaling in angiogenesis. <u>Anal Chem. 87 (6): 3239-46.</u>
- 12. Hilbe W *et al.* (2004) CD133 positive endothelial progenitor cells contribute to the tumour vasculature in non-small cell lung cancer. <u>J Clin Pathol. 57 (9): 965-9.</u>
- 13. Yi, T. *et al.* (2015) Manufacture of Clinical-Grade Human Clonal Mesenchymal Stem Cell Products from Single Colony Forming Unit-Derived Colonies Based on the Subfractionation Culturing Method. Tissue Eng Part C Methods. 21 (12): 1251-62.
- 14. Palakkan, A.A. *et al.* (2015) Polarisation and functional characterisation of hepatocytes derived from human embryonic and mesenchymal stem cells. <u>Biomed Rep. 3 (5):</u> 626-636.
- 15. Newey SE *et al.* (2014) The hematopoietic chemokine CXCL12 promotes integration of human endothelial colony forming cell-derived cells into immature vessel networks. Stem Cells Dev. 23 (22): 2730-43.
- 16. Fabre-Mersseman V *et al.* (2011) CD4<sup>+</sup> recent thymic emigrants are infected by HIV in vivo, implication for pathogenesis. <u>AIDS. 25 (9): 1153-62.</u>
- 17. Patten PE *et al.* (2008) CD38 expression in chronic lymphocytic leukemia is regulated by the tumor microenvironment. <u>Blood. 111 (10): 5173-81.</u>
- 18. Katz SC *et al.* (2004) Liver sinusoidal endothelial cells are insufficient to activate T cells. J Immunol. 173 (1): 230-5.
- 19. Pfisterer K *et al.* (2015) CD90(+) human dermal stromal cells are potent inducers of FoxP3(+) regulatory T cells. J Invest Dermatol. 135 (1): 130-41.
- 20. Hale, S.J. *et al.* (2015) CXCR2 modulates bone marrow vascular repair and haematopoietic recovery post-transplant. Br J Haematol. 169 (4): 552-64.
- 21. Muthana, M. *et al.* (2015) Directing cell therapy to anatomic target sites in vivo with magnetic resonance targeting. <u>Nat Commun. 6: 8009.</u>
- 22. Schuster, C. *et al.* (2015) Development of Blood and Lymphatic Endothelial Cells in Embryonic and Fetal Human Skin. <u>Am J Pathol. 185 (9): 2563-74.</u>
- 23. Somers, E. *et al.* (2016) Vascular Defects and Spinal Cord Hypoxia in Spinal Muscular Atrophy. <u>Ann Neurol. 79 (2): 217-30.</u>
- 24. Soh, B.S. *et al.* (2016) Endothelin-1 supports clonal derivation and expansion of cardiovascular progenitors derived from human embryonic stem cells. <u>Nat Commun. 7:</u> 10774.
- 25. GarikipatiV, N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. <u>PLoS One. 13 (2): e0192244.</u>
- 26. Duque, J.C. *et al.* (2019) Vascularization of the arteriovenous fistula wall and association with maturation outcomes. <u>J Vasc Access.: 1129729819863584. [Epub ahead of print]</u>

**Storage** 

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and

should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

| Guarantee                        | 12 months from date of despatch  |
|----------------------------------|--|
| Health And Safety<br>Information | Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1738F">https://www.bio-rad-antibodies.com/SDS/MCA1738F</a> 10041 |
| Regulatory                       | For research purposes only   |

# **Related Products**

## **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

## **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365643:200529'

## Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint