

Datasheet: MCA1742 BATCH NUMBER 154454

Description:	MOUSE ANTI HUMAN CD227
Specificity:	CD227
Other names:	CA 15-3, MUCIN 1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	C595 (NCRC48)
Isotype:	lgG3
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			
Immunohistology - Frozen	•			
Immunohistology - Paraffin (1)	•			
ELISA	•			1/100 - 1/1000
Immunoprecipitation				
Western Blotting	•			
Immunofluorescence				

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)This product requires heat mediated antigen retrieval prior to staining of paraffin embedded material. Citrate buffer pH6.0 is recommended for this purpose.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein A supernatant	A from tissue culture

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Urinary MUC-1 mucin.
External Database Links	UniProt: P15941 Related reagents Entrez Gene: 4582 MUC1 Related reagents
Synonyms	PUM
RRID	
KKID	AB_322211
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of a mouse myeloma cell line.
Specificity	Mouse anti Human CD227 antibody, clone C595 (NCRC48) recognizes CD227, also known as mucin 1 which is a breast cancer associated mucin encoded by the Muc-1 gene. Mucins are a family of high molecular weight, heavily glycosylated proteins (glycoconjugates) produced by many epithelial tissues in vertebrates. CD227 is expressed on most secretory epithelium, including mammary gland and some hematopoietic cells. This protein is overexpressed abundantly in >90% breast carcinomas and metastases. Mouse anti Human CD227 antibody, clone C595 recognizes the peptide epitope ARG-PRO-ALA-PRO within the protein core of the mucin.
Histology Positive Control Tissue	Breast carcinoma
References	 Price, M.R. <i>et al.</i> (1990) Immunological and structural features of the protein core of human polymorphic epithelial mucin. Mol Immunol. 27 (8): 795-802. Price, M.R. <i>et al.</i> (1990) C595a monoclonal antibody against the protein core of human urinary epithelial mucin commonly expressed in breast carcinomas. Br J Cancer. 61 (5): 681-6. Price, M.R. <i>et al.</i> (1991) Purification of anti-epithelial mucin monoclonal antibodies by epitope affinity chromatography. J Immunol Methods. 139 (1): 83-90. Denton, G. <i>et al.</i> (1995) Primary sequence determination and molecular modelling of the variable region of an antiMUC1 mucin monoclonal antibody. Eur J Cancer. 31A (2): 214-21.

- 5. Denton, G. *et al.* (1997) Production and characterization of a recombinant anti-MUC1 scFv reactive with human carcinomas. Br J Cancer. 76 (5): 614-21.
- 6. Murray, A. *et al* (2001) Production and characterization of 188Re-C595 antibody for radioimmunotherapy of transitional cell bladder cancer. <u>J Nucl Med. 42: 726-32.</u>
- 7. Hisatsune, A. *et al.* (2011) Anti-MUC1 antibody inhibits EGF receptor signaling in cancer cells. Biochem Biophys Res Commun. 405: 377-81.
- 8. Ragupathi, G. *et al.* (2005) Antibodies against tumor cell glycolipids and proteins, but not mucins, mediate complement-dependent cytotoxicity. <u>J Immunol</u>. 174: 5706-12.
- 9. Huang, D.M. *et al.* (2004) Modulation of anti-adhesion molecule MUC-1 is associated with arctiin-induced growth inhibition in PC-3 cells. <u>Prostate. 2004 May 15;59(3): 260-7.</u>

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1742 10040
Regulatory	For research purposes only

Related Products

North & South Tel: +1 800 265 7376

Fax: +1 919 878 3751

America

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)

Rabbit Anti Mouse IgG (STAR12...)

RPE

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Goat Anti Mouse IgG (STAR76...) RPE

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

Worldwide

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

Tel: +44 (0)1865 852 700 **Europe** Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M365651:200529'

Fax: +44 (0)1865 852 739

Printed on 25 Mar 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint