

Datasheet: MCA1744F

Description:	MOUSE ANTI HUMAN CD66e:FITC
Specificity:	CD66e
Other names:	CEA
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	C365D3 (NCRC23)
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% sodium azide (NaN ₃)		
Stabilisers	1% bovine serum albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		

External Database
Links

UniProt:

[P06731](#)

[Related reagents](#)

Entrez Gene:

[1048](#)

CEACAM5

[Related reagents](#)

Synonyms

CEA

RRID

AB_323914

Fusion Partners

Spleen cells from immunised BALB/c mice were fused with cells of the mouse P3NSI myeloma cell line.

Specificity

Mouse anti Human CD66e antibody, clone C365D3 (NCRC23) recognizes human Carcinoembryonic antigen-related cell adhesion molecule 5, also known as CD66e, carcinoembryonic antigen, Meconium antigen 100, CEA or CEACAM5. CD66e is a 702 amino acid ~77 kDa GPI anchored membrane protein containing 7 [Ig-like domains](#). Mouse anti Human CD66e antibody, clone C365D3 does not cross-react with normal cross-reacting antigen (CD66c), or with biliary glycoprotein 1 (CD66a) as indicated by binding assays ([Price 1988](#), note: in this study Mouse anti Human CD66e antibody, clone C365D3 is designated as clone 6 (from author)).

Flow Cytometry

Use 10µl of the suggested working dilution to label 10⁶ cells in 100µl

References

1. Seth, J. *et al.* (1988) Carcinoembryonic antigen. [Lancet. 1 \(8599\): 1399.](#)
2. Chao, A. *et al.* (2006) Molecular characterization of adenocarcinoma and squamous carcinoma of the uterine cervix using microarray analysis of gene expression. [Int J Cancer. 119: 91-8.](#)
3. Stern-Ginossar, N. *et al.* (2007) Intercellular transfer of carcinoembryonic antigen from tumor cells to NK cells. [J Immunol. 179 \(7\): 4424-34.](#)
4. Kalinina, T. *et al.* (2010) Establishment and characterization of a new human pancreatic adenocarcinoma cell line with high metastatic potential to the lung. [BMC Cancer.10: 295.](#)
5. Soucek, K. *et al.* (2010) Fetal colon cell line FHC exhibits tumorigenic phenotype, complex karyotype, and TP53 gene mutation. [Cancer Genet Cytogenet. 197: 107-16.](#)
6. Ferro, F. *et al.* (2011) Adipose tissue-derived stem cell in vitro differentiation in a three-dimensional dental bud structure. [Am J Pathol.178: 2299-310.](#)
7. Dallas, M.R. *et al.* (2012) Divergent roles of CD44 and carcinoembryonic antigen in colon cancer metastasis. [FASEB J. 226: 2648-56.](#)
8. Domenis, R. *et al.* (2015) Adipose tissue derived stem cells: in vitro and in vivo analysis of a standard and three commercially available cell-assisted lipotransfer techniques. [Stem Cell Res Ther. 6: 2.](#)
9. Wicklein, D. *et al.* (2018) CEACAM1 promotes melanoma metastasis and is involved in the regulation of the EMT associated gene network in melanoma cells. [Sci Rep. 8 \(1\): 11893.](#)
10. Caponnetto, F. *et al.* (2020) Human Adipose-Derived Stem Cells in Madelung's Disease: Morphological and Functional Characterization. [Cells: 10 \(1\): 44.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1744F 10041
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)
[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M411304:221102'

Printed on 24 May 2025