

## Datasheet: MCA1749GA

<b>Description:</b>	MOUSE ANTI PIG CD4 ALPHA
<b>Specificity:</b>	CD4 ALPHA
<b>Other names:</b>	CD4
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	MIL17
<b>Isotype:</b>	IgG2b
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/200
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.**

<b>Target Species</b>	Pig
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline

<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Leucocytes isolated from porcine gut lamina propria.
<b>Specificity</b>	<p><b>Mouse anti Porcine CD4 alpha, clone MIL17</b> recognizes a ~55 kDa porcine homologue to the human CD4 antigen found on the surface of helper-T cells. MIL-17 was confirmed as a member of the CD4 alpha cluster at the 'Third International Workshop on Swine Leukocyte Differentiation Antigens' (<a href="#">Haverson et al. 2001</a>). Porcine CD4 is a type 1 trans-membrane member of the immunoglobulin superfamily.</p> <p>Pigs appear unusual amongst mammalian species as they appear to have four populations of resting T lymphocytes. In addition to the two populations of mutually exclusive CD4+/CD8- and CD4-/CD8+ lymphocytes, they also appear to have significant populations of CD4-/CD8- and CD4+/CD8+ cells. Lymphoblasts with a double positive phenotype have been described in other species but this is not the case for mature T lymphocytic cells (<a href="#">Saalmuller et al. 1987</a>)</p> <p>Mouse anti Pig CD4 alpha, clone MIL17 stains a population of cells with characteristic lymphocyte morphology in immunohistochemistry (Inman <i>et al.</i> 2010).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul
<b>References</b>	<ol style="list-style-type: none"> <li>1. Saalmüller A <i>et al.</i> (2001) Summary of workshop findings for porcine T-lymphocyte-specific monoclonal antibodies. <a href="#">Vet Immunol Immunopathol. 80 (1-2): 35-52.</a></li> <li>2. Castellano, G. <i>et al.</i> (2010) Therapeutic targeting of classical and lectin pathways of complement protects from ischemia-reperfusion-induced renal damage. <a href="#">Am J Pathol. 176: 1648-59.</a></li> <li>3. Inman, C.F. <i>et al.</i> (2010) Dendritic cells interact with CD4 T cells in intestinal mucosa. <a href="#">J Leukoc Biol. 88 (3): 571-8.</a></li> <li>4. Kick AR <i>et al.</i> (2011) Evaluation of peripheral lymphocytes after weaning and vaccination for <i>Mycoplasma hyopneumoniae</i>. <a href="#">Res Vet Sci. 91 (3): e68-72.</a></li> <li>5. Kick, A.R. <i>et al.</i> (2012) Effects of stress associated with weaning on the adaptive immune system in pigs. <a href="#">J Anim Sci. 90: 649-56.</a></li> <li>6. Goujon, J.M. <i>et al.</i> (2000) Influence of cold-storage conditions on renal function of autotransplanted large pig kidneys. <a href="#">Kidney Int. 58: 838-50.</a></li> <li>7. Tambuyzer BR <i>et al.</i> (2012) Osteopontin alters the functional profile of porcine microglia <i>in vitro</i>. <a href="#">Cell Biol Int. 36 (12): 1233-8.</a></li> <li>8. Tuchscherer, M. <i>et al.</i> (2012) Effects of inadequate maternal dietary protein:carbohydrate ratios during pregnancy on offspring immunity in pigs. <a href="#">BMC Vet Res. 8: 232.</a></li> <li>9. Cao, D. <i>et al.</i> (2010) Synthetic innate defence regulator peptide enhances in vivo immunostimulatory effects of CpG-ODN in newborn piglets. <a href="#">Vaccine. 28: 6006-13.</a></li> </ol>

10. Clapperton, M. *et al.* (2005) Associations of weight gain and food intake with leukocyte sub-sets in Large White pigs [Livestock Production Science 96: 249-60](#)
11. Clapperton, M. *et al.* (2005) Innate immune traits differ between Meishan and Large White pigs. [Vet Immunol Immunopathol. 104: 131-44.](#)
12. Clapperton, M. *et al.* (2008) Pig peripheral blood mononuclear leucocyte subsets are heritable and genetically correlated with performance. [Animal. 2: 1575-84.](#)
13. Faure, J.P. *et al.* (2002) Polyethylene glycol reduces early and long-term cold ischemia-reperfusion and renal medulla injury. [J Pharmacol Exp Ther. 2002 Sep;302\(3\):861-70.](#)
14. Faure, J.P. *et al.* (2004) Evidence for protective roles of polyethylene glycol plus high sodium solution and trimetazidine against consequences of renal medulla ischaemia during cold preservation and reperfusion in a pig kidney model. [Nephrol Dial Transplant. 19: 1742-51.](#)
15. Inman, C.F. *et al.* (2012) Neonatal colonisation expands a specific intestinal antigen-presenting cell subset prior to CD4 T-cell expansion, without altering T-cell repertoire. [PLoS One. 7\(3\): e33707.](#)
16. Kick, A.R. *et al.* (2012) Effects of stress associated with weaning on the adaptive immune system in pigs. [J Anim Sci. 90: 649-56.](#)
17. Langerhuus, S.N. *et al.* (2010) Brief report: biomarkers of aortic vascular prosthetic graft infection in a porcine model with *Staphylococcus aureus*. [Eur J Clin Microbiol Infect Dis. 29: 1453-6.](#)
18. Lu, X. *et al.* (2012) Genome-wide association study for T lymphocyte subpopulations in swine. [BMC Genomics. 13: 488.](#)
19. Monroy-Salazar, H.G. *et al.* (2012) Effects of a live yeast dietary supplement on fecal coliform counts and on peripheral blood CD4+ and CD8+ lymphocyte subpopulations in nursery pigs. [J Swine Health Prod 20: 276-282.](#)
20. Shi, K. *et al.* (2008) Changes in peripheral blood leukocyte subpopulations in piglets co-infected experimentally with porcine reproductive and respiratory syndrome virus and porcine circovirus type 2. [Vet Microbiol. 129: 367-77.](#)
21. Spreeuwenberg, M.A. *et al.* (2001) Small intestine epithelial barrier function is compromised in pigs with low feed intake at weaning. [J Nutr. 131: 1520-7.](#)
22. Tambuyzer, B.R. *et al.* (2012) Osteopontin alters the functional profile of porcine microglia in vitro. [Cell Biol Int. 36: 1233-8.](#)
23. Zelnickova, P. *et al.* (2007) Intracellular cytokine detection by flow cytometry in pigs: fixation, permeabilization and cell surface staining. [J Immunol Methods. 327: 18-29.](#)
24. Kvist, P.H. *et al.* (2010) Effect of subcutaneous glucose sensor implantation on skin mRNA expression in pigs. [Diabetes Technol Ther. 12: 791-9.](#)
25. Lefevre, E.A. *et al.* (2012) Immune responses in pigs vaccinated with adjuvanted and non-adjuvanted A(H1N1)pdm/09 influenza vaccines used in human immunization programmes. [PLoS One. 7\(3\): e32400.](#)
26. Akershoek, J.J. *et al.* (2016) Cell therapy for full-thickness wounds: are fetal dermal cells a potential source? [Cell Tissue Res. 364 \(1\): 83-94.](#)
27. Liu J *et al.* (2016) The Role of Porcine Monocyte Derived Dendritic Cells (MoDC) in the Inflammation Storm Caused by *Streptococcus suis* Serotype 2 Infection. [PLoS One. 11 \(3\): e0151256.](#)
28. Liermann, W. *et al.* (2017) Effects of two commercial diets and technical feed treatment on stomach lesions and immune system of fattening pigs. [J Anim Physiol Anim](#)

[Nutr \(Berl\). 101 \(5\): e414-26.](#)

29. Gardner, D.S. *et al.* (2016) Remote effects of acute kidney injury in a porcine model. [Am J Physiol Renal Physiol. 310 \(4\): F259-71.](#)

30. Hemmink, J.D. *et al.* (2016) Distinct immune responses and virus shedding in pigs following aerosol, intra-nasal and contact infection with pandemic swine influenza A virus, A(H1N1)09. [Vet Res. 47 \(1\): 103.](#)

31. Dąbrowski, M. *et al.* (2017) The Effect of Deoxynivalenol on Selected Populations of Immunocompetent Cells in Porcine Blood-A Preliminary Study. [Molecules. 22 \(5\)Apr 26 \[Epub ahead of print\].](#)

32. Hsu, W.T. *et al.* (2013) Prostaglandin E2 potentiates mesenchymal stem cell-induced IL-10+IFN- $\gamma$ +CD4+ regulatory T cells to control transplant arteriosclerosis. [J Immunol. 190 \(5\): 2372-80.](#)

33. Matsubara, T. *et al.* (2015) Identification of a CD4 variant in Microminipigs not detectable with available anti-CD4 monoclonal antibodies. [Vet Immunol Immunopathol. 168 \(3-4\): 176-83.](#)

34. Hu, Z. *et al.* (2019) Genomic variant in porcine TNFRSF1A gene and its effects on TNF signaling pathway *in vitro*. [Gene. 700: 105-9.](#)

35. Fogle, J.E. *et al.* (2019) Antibiotic Therapy Does Not Alter the Humoral Response to Vaccination for Porcine Circovirus 2 in Weaned Pigs. [Vet Sci. 6\(2\): 51.](#)

36. Forner, R. *et al.* (2021) Distribution difference of colostrum-derived B and T cells subsets in gilts and sows. [PLoS One. 16 \(5\): e0249366.](#)

37. Christoforidou, Z. *et al.* (2019) Sexual Dimorphism in Immune Development and in Response to Nutritional Intervention in Neonatal Piglets. [Front Immunol. 10: 2705.](#)

38. López, E. *et al.* (2019) Identification of very early inflammatory markers in a porcine myocardial infarction model. [BMC Vet Res. 15 \(1\): 91.](#)

39. Liu, K.Y. *et al.* (2021) Fallopian tube stem cell medium of porcine and bovine: *In vitro* regenerative effect on maturation and parthenogenesis of porcine oocytes. [Res Vet Sci. 140: 83-90.](#)

40. Giese, I.M. *et al.* (2020) Chronic Hyperglycemia Drives Functional Impairment of Lymphocytes in Diabetic *INS*<sup>C94Y</sup> Transgenic Pigs. [Front Immunol. 11: 607473.](#)

41. Nielsen, O.L. *et al.* (2021) A porcine model of subcutaneous *Staphylococcus aureus* infection: a pilot study. [APMIS. Mar 01 \[Epub ahead of print\].](#)

---

**Further Reading** 1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR8...)	<a href="#">DyLight®800</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL \(MCA691\)](#)

### Recommended Useful Reagents

[MOUSE ANTI PIG CD45 \(MCA1222GA\)](#)

[MOUSE ANTI PIG wCD8 ALPHA \(MCA1223GA\)](#)

[RAT ANTI HUMAN CD3 \(MCA1477\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M383150:210513'

Printed on 07 Jan 2022