

Datasheet: MCA1751F

BATCH NUMBER 1801

Description:	MOUSE ANTI PIG CD45RA:FITC
Specificity:	CD45RA
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	MIL13
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Pig		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1.0% Bovine Serum Albumin		
Approx. Protein	IgG concentration 0.1 mg/ml		

Concentrations

Immunogen Cells isolated from porcine mesenteric lymph node

RRID AB_323348

Specificity **Mouse anti Pig CD45RA, clone MIL13**, recognizes an epitope contained in the portion of porcine CD45 encoded by exon A, CD45RA ([Lunney et al. 2007](#)).

Mouse anti pig CD45RA, clone MIL13 recognizes both the 210 kDa RA CD45 isoform and the 226 kDa RAC isoform ([Zuckermann et al. 2001](#)). Clone MIL13 does not recognize the CD45RC or CD45RO isoforms.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

- References**
1. Bozić F *et al.* (2002) Recruitment of intestinal CD45RA+ and CD45RC+ cells induced by a candidate oral vaccine against porcine post-weaning colibacillosis. [Vet Immunol Immunopathol. 86 \(3-4\): 137-46.](#)
 2. Terzic, S. *et al.* (2002) Immunophenotyping of leukocyte subsets in peripheral blood and palatine tonsils of prefattening pigs. [Vet Res Commun. 26: 273-83.](#)
 3. Pakkanen, T.M. *et al.* (2000) Periadventitial lacZ gene transfer to pig carotid arteries using a biodegradable collagen collar or a wrap of collagen sheet with adenoviruses and plasmid-liposome complexes. [J Gene Med. 2: 52-60.](#)
 4. Schierack, P. *et al.* (2009) Effects of *Bacillus cereus* var. *toyoi* on immune parameters of pregnant sows. [Vet Immunol Immunopathol. 127: 26-37.](#)
 5. Suzuki, S. *et al.* (2016) Generation and characterization of RAG2 knockout pigs as animal model for severe combined immunodeficiency. [Vet Immunol Immunopathol. 178: 37-49.](#)
 6. López, E. *et al.* (2019) Identification of very early inflammatory markers in a porcine myocardial infarction model. [BMC Vet Res. 15 \(1\): 91.](#)
 7. Li, K. *et al.* (2019) Generation of porcine monoclonal antibodies based on single cell technologies. [Vet Immunol Immunopathol. 215: 109913.](#)
-

Further Reading 1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

Storage Store at +4°C for one month or at -20°C for longer.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1751F>

10041

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M365676:200529'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)