



Datasheet: MCA1757A647

**BATCH NUMBER 170870**

<b>Description:</b>	MOUSE ANTI HUMAN MYELOPEROXIDASE:Alexa Fluor® 647
<b>Specificity:</b>	MYELOPEROXIDASE
<b>Format:</b>	ALEXA FLUOR® 647
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	2C7
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS/1ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code [BUF09](#)) is recommended for this purpose.**

### Target Species

Human

### Species Cross Reactivity

Reacts with: Dog

Does not react with:Rat

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG conjugated to Alexa Fluor® 647 - liquid

### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
Alexa Fluor®647	650	665

### Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture

supernatant

---

**Buffer Solution** Phosphate buffered saline

---

**Preservative** 0.09% sodium azide (NaN<sub>3</sub>)  
**Stabilisers** 1% bovine serum albumin

---

**Approx. Protein Concentrations** IgG concentration 0.05 mg/ml

---

**Immunogen** Human myeloperoxidase

---

**External Database Links**

**UniProt:**

[P05164](#) [Related reagents](#)

**Entrez Gene:**

[4353](#) MPO [Related reagents](#)

---

**Fusion Partners** Spleen cells from immunized mice were fused with cells of the mouse X63 AG8-653 myeloma cell line

---

**Specificity** **Mouse anti Human myeloperoxidase antibody, clone 2C7** recognizes human myeloperoxidase (MPO). MPO is an important component of azurophilic granules in neutrophils, being involved in microbicidal processes. The protein is a multimer of 2 heavy chains (55 kDa) and two light chains (15 kDa), the heavy chains being linked by a disulphide bond.

Mouse anti Human Myeloperoxidase antibody, clone 2C7 recognizes native MPO in Western blots, and the heavy chain following boiling of the sample. Mouse anti Human Myeloperoxidase antibody, clone 2C7 also recognizes recombinant MPO in western blots and weakly in ELISA.

Mouse anti Human myeloperoxidase antibody, clone 2C7 may be of value in the study of myeloid cells and myeloid leukaemias by flow cytometry following cell permeabilization. Mouse anti Human myeloperoxidase antibody, clone 2C7 did not recognize rat MPO by ELISA ([Patry et al. 2003](#)).

---

**Flow Cytometry** Use 10µl of the suggested working dilution to label 10<sup>6</sup> cells in 100µl

---

**References**

1. Patry, Y.C. *et al.* (2003) Difference in antigenic determinant profiles between human and rat myeloperoxidase. [Clin Exp Immunol. 132 \(3\): 505-8.](#)
2. Sloane, A.J. *et al.* (2005) Proteomic analysis of sputum from adults and children with cystic fibrosis and from control subjects. [Am J Respir Crit Care Med. 172: 1416-26.](#)
3. Villiers, E. *et al.* (2006) Identification of acute myeloid leukemia in dogs using flow cytometry with myeloperoxidase, MAC387, and a canine neutrophil-specific antibody. [Vet Clin Pathol. 35 \(1\): 55-71.](#)
4. Villiers, E. *et al.* (2006) Identification of acute myeloid leukemia in dogs using flow

cytometry with myeloperoxidase, MAC387, and a canine neutrophil-specific antibody. [Vet Clin Pathol. 35 \(1\): 55-71.](#)

5. Luo, B. *et al.* (2013) Immunopathology features of chronic rhinosinusitis in high-altitude dwelling Tibetans. [Allergy Rhinol \(Providence\). 4: e69-76.](#)
6. Gelain, M.E. *et al.* (2014) CD44 in canine leukemia: analysis of mRNA and protein expression in peripheral blood. [Vet Immunol Immunopathol. 159 \(1-2\): 91-6.](#)
7. Behnen, M. *et al.* (2014) Immobilized immune complexes induce neutrophil extracellular trap release by human neutrophil granulocytes via FcγRIIIB and Mac-1. [J Immunol. 193 \(4\): 1954-65.](#)
8. Novacco, M. *et al.* (2016) Prognostic factors in canine acute leukaemias: a retrospective study. [Vet Comp Oncol. 14 \(4\): 409-16.](#)
9. Wang, H. *et al.* (2016) Circulating Level of Neutrophil Extracellular Traps Is Not a Useful Biomarker for Assessing Disease Activity in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. [PLoS One. 11 \(2\): e0148197.](#)
10. Rai, A.K. *et al.* (2017) Exonal switch down-regulates the expression of CD5 on blasts of acute T cell leukaemia. [Clin Exp Immunol. 190 \(3\): 340-350.](#)
11. Hoppenbrouwers, T. *et al.* (2018) Neutrophil Extracellular Traps in Children With Meningococcal Sepsis. [Pediatr Crit Care Med. 19 \(6\): e286-e291.](#)
12. Helseth, R. *et al.* (2019) Glucose associated NETosis in patients with ST-elevation myocardial infarction: an observational study. [BMC Cardiovasc Disord. 19 \(1\): 221.](#)
13. Manfredi, A.A. *et al.* (2021) Platelet phagocytosis via PSGL1 and accumulation of microparticles in systemic sclerosis. [Arthritis Rheumatol. Jul 19 \[Epub ahead of print\].](#)
14. de Moraes Mazetto, B. *et al.* (2022) Association between neutrophil extracellular traps (NETs) and thrombosis in antiphospholipid syndrome [Thrombosis Res. 214: 132-7.](#)
15. Robinson, H. *et al.* (2022) The effect of expressive writing on wound healing: Immunohistochemistry analysis of skin tissue two weeks after punch biopsy wounding. [J Psychosom Res. 161: 110987.](#)
16. Zhu, D. *et al.* (2024) Neutrophil activation biomarker pentraxin 3 for diagnosis and monitoring of macrophage activation syndrome occurrence in adult-onset Still's disease. [J Autoimmun. 144: 103182.](#)
17. Rogato, F. *et al.* (2024) Leukemia cutis as a prominent clinical sign in a dog with acute myeloid leukemia. [Vet Clin Pathol. 53 \(4\): 448-57.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

---

**Guarantee** 12 months from date of despatch

---

**Acknowledgements** This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchased product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening

services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or [outlicensing@thermofisher.com](mailto:outlicensing@thermofisher.com)

---

**Health And Safety Information**      Material Safety Datasheet documentation #10041 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA1757A647>

---

**Regulatory**                      For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA928A647\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**Product inquiries:** [www.bio-rad-antibodies.com/technical-support](http://www.bio-rad-antibodies.com/technical-support)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](http://bio-rad-antibodies.com/datasheets)  
'M438127:250327'

**Printed on 25 Feb 2026**

---

© 2026 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)