

Datasheet: MCA1757PE

Description:	MOUSE ANTI HUMAN MYELOPEROXIDASE:RPE
Specificity:	MYELOPEROXIDASE
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	2C7
Isotype:	lgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	•			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code <u>BUF09</u>) is recommended for this purpose.

Target Species	Human			
Species Cross	Reacts with: Dog			
Reactivity	Does not react with	:Rat		
	reactivity is derived	from testing within our la	ons may vary between s aboratories, peer-reviewe ors. Please refer to refere	ed publications or
Product Form	Purified IgG conjuga	ated to R. Phycoerythrin	(RPE) - lyophilized	
Reconstitution	Reconstitute with 1	ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	RPE 488nm laser	496	578	

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃) 1% bovine serum albumin 5% sucrose
Immunogen	Human myeloperoxidase
External Database Links	UniProt: P05164 Related reagents Entrez Gene: 4353 MPO Related reagents
RRID	AB_322880
Fusion Partners	Spleen cells from immunized mice were fused with cells of the mouse X63 AG8-653 myeloma cell line
Specificity	Mouse anti Human myeloperoxidase antibody, clone 2C7 recognizes human myeloperoxidase (MPO). MPO is an important component of azurophilic granules in neutrophils, being involved in microbicidal processes. The protein is a multimer of 2 heavy chains (55 kDa) and two light chains (15 kDa), the heavy chains being linked by a disulphide bond.
	Mouse anti Human Myeloperoxidase antibody, clone 2C7 recognizes native MPO in Western blots, and the heavy chain following boiling of the sample. Mouse anti Human Myeloperoxidase antibody, clone 2C7 also recognizes recombinant MPO in western blots and weakly in ELISA.
	Mouse anti Human myeloperoxidase antibody, clone 2C7 may be of value in the study of myeloid cells and myeloid leukaemias by flow cytometry following cell permeabilization. Mouse anti Human myeloperoxidase antibody, clone 2C7 did not recognize rat MPO by ELISA (<u>Patry et al. 2003</u>).
Flow Cytometry	Use 10μl of the suggested working dilution to label 10 ⁶ cells in 100μl
References	 Patry, Y.C. <i>et al.</i> (2003) Difference in antigenic determinant profiles between human and rat myeloperoxidase. <u>Clin Exp Immunol. 132 (3): 505-8.</u> Sloane, A.J. <i>et al.</i> (2005) Proteomic analysis of sputum from adults and children with cystic fibrosis and from control subjects. <u>Am J Respir Crit Care Med. 172: 1416-26.</u> Villiers, E. <i>et al.</i> (2006) Identification of acute myeloid leukemia in dogs using flow cytometry with myeloperoxidase, MAC387, and a canine neutrophil-specific antibody. <u>Vet</u>

Clin Pathol. 35 (1): 55-71.

- 4. Villiers, E. *et al.* (2006) Identification of acute myeloid leukemia in dogs using flow cytometry with myeloperoxidase, MAC387, and a canine neutrophil-specific antibody. <u>Vet Clin Pathol. 35 (1): 55-71</u>.
- 5. Luo, B. *et al.* (2013) Immunopathology features of chronic rhinosinusitis in high-altitude dwelling Tibetans. <u>Allergy Rhinol (Providence)</u>. 4: e69-76.
- 6. Gelain, M.E. *et al.* (2014) CD44 in canine leukemia: analysis of mRNA and protein expression in peripheral blood. <u>Vet Immunol Immunopathol.</u> 159 (1-2): 91-6.
- 7. Behnen, M. *et al.* (2014) Immobilized immune complexes induce neutrophil extracellular trap release by human neutrophil granulocytes via FcγRIIB and Mac-1. <u>J Immunol. 193</u> (4): 1954-65.
- 8. Novacco, M. *et al.* (2016) Prognostic factors in canine acute leukaemias: a retrospective study. Vet Comp Oncol. 14 (4): 409-16.
- 9. Wang, H. *et al.* (2016) Circulating Level of Neutrophil Extracellular Traps Is Not a Useful Biomarker for Assessing Disease Activity in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. PLoS One. 11 (2): e0148197.
- 10. Rai, A.K. *et al.* (2017) Exonal switch down-regulates the expression of CD5 on blasts of acute T cell leukaemia. Clin Exp Immunol. 190 (3): 340-350.
- 11. Hoppenbrouwers, T. *et al.* (2018) Neutrophil Extracellular Traps in Children With Meningococcal Sepsis. <u>Pediatr Crit Care Med. 19 (6): e286-e291.</u>
- 12. Helseth, R. *et al.* (2019) Glucose associated NETosis in patients with ST-elevation myocardial infarction: an observational study. <u>BMC Cardiovasc Disord.</u> 19 (1): 221.
- 13. Manfredi, A.A. *et al.* (2021) Platelet phagocytosis via PSGL1 and accumulation of microparticles in systemic sclerosis. Arthritis Rheumatol. Jul 19 [Epub ahead of print].
- 14. de Moraes Mazetto, B. *et al.* (2022) Association between neutrophil extracellular traps (NETs) and thrombosis in antiphospholipid syndrome <u>Thrombosis Res. 214: 132-7.</u>
- 15. Robinson, H. *et al.* (2022) The effect of expressive writing on wound healing: Immunohistochemistry analysis of skin tissue two weeks after punch biopsy wounding. <u>J. Psychosom Res.</u> 161: 110987.
- 16. Zhu, D. *et al.* (2024) Neutrophil activation biomarker pentraxin 3 for diagnosis and monitoring of macrophage activation syndrome occurrence in adult-onset Still's disease. <u>J</u> Autoimmun. 144: 103182.
- 17. Rogato, F. *et al.* (2024) Leukemia cutis as a prominent clinical sign in a dog with acute myeloid leukemia. <u>Vet Clin Pathol. 53 (4): 448-57.</u>

Storage

This product is shipped at ambient temperature.

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA1757PE 20487

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M440251:250523'

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint