

## Datasheet: MCA1757T

<b>Description:</b>	MOUSE ANTI HUMAN MYELOPEROXIDASE
<b>Specificity:</b>	MYELOPEROXIDASE
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	2C7
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	20 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			1/50 - 1/100
Immunohistology - Frozen	▪			1/1000 - 1/5000
Immunohistology - Paraffin	▪			1/500 - 1/1000
ELISA	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code [BUF09](#)) is recommended for this purpose.**

### Target Species

Human

### Species Cross Reactivity

Reacts with: Dog

Does not react with: Rat

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG - liquid

### Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% sodium azide (NaN <sub>3</sub> )
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Human myeloperoxidase
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P05164</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">4353</a>    MPO    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_2146465
<b>Fusion Partners</b>	Spleen cells from immunized mice were fused with cells of the mouse X63 AG8-653 myeloma cell line
<b>Specificity</b>	<p><b>Mouse anti Human myeloperoxidase antibody, clone 2C7</b> recognizes human myeloperoxidase (MPO). MPO is an important component of azurophilic granules in neutrophils, being involved in microbicidal processes. The protein is a multimer of 2 heavy chains (55 kDa) and two light chains (15 kDa), the heavy chains being linked by a disulphide bond.</p> <p>Mouse anti Human Myeloperoxidase antibody, clone 2C7 recognizes native MPO in Western blots, and the heavy chain following boiling of the sample. Mouse anti Human Myeloperoxidase antibody, clone 2C7 also recognizes recombinant MPO in western blots and weakly in ELISA.</p> <p>Mouse anti Human myeloperoxidase antibody, clone 2C7 may be of value in the study of myeloid cells and myeloid leukaemias by flow cytometry following cell permeabilization. Mouse anti Human myeloperoxidase antibody, clone 2C7 did not recognize rat MPO by ELISA (<a href="#">Patry et al. 2003</a>).</p>
<b>Flow Cytometry</b>	Use 10µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl
<b>Histology Positive Control Tissue</b>	Human bone marrow
<b>References</b>	<ol style="list-style-type: none"> <li>1. Patry, Y.C. <i>et al.</i> (2003) Difference in antigenic determinant profiles between human and rat myeloperoxidase. <a href="#">Clin Exp Immunol. 132 (3): 505-8.</a></li> <li>2. Sloane, A.J. <i>et al.</i> (2005) Proteomic analysis of sputum from adults and children with cystic fibrosis and from control subjects. <a href="#">Am J Respir Crit Care Med. 172: 1416-26.</a></li> </ol>

3. Villiers, E. *et al.* (2006) Identification of acute myeloid leukemia in dogs using flow cytometry with myeloperoxidase, MAC387, and a canine neutrophil-specific antibody. [Vet Clin Pathol. 35 \(1\): 55-71.](#)
4. Villiers, E. *et al.* (2006) Identification of acute myeloid leukemia in dogs using flow cytometry with myeloperoxidase, MAC387, and a canine neutrophil-specific antibody. [Vet Clin Pathol. 35 \(1\): 55-71.](#)
5. Luo, B. *et al.* (2013) Immunopathology features of chronic rhinosinusitis in high-altitude dwelling Tibetans. [Allergy Rhinol \(Providence\). 4: e69-76.](#)
6. Gelain, M.E. *et al.* (2014) CD44 in canine leukemia: analysis of mRNA and protein expression in peripheral blood. [Vet Immunol Immunopathol. 159 \(1-2\): 91-6.](#)
7. Behnen, M. *et al.* (2014) Immobilized immune complexes induce neutrophil extracellular trap release by human neutrophil granulocytes via FcγRIIIB and Mac-1. [J Immunol. 193 \(4\): 1954-65.](#)
8. Novacco, M. *et al.* (2016) Prognostic factors in canine acute leukaemias: a retrospective study. [Vet Comp Oncol. 14 \(4\): 409-16.](#)
9. Wang, H. *et al.* (2016) Circulating Level of Neutrophil Extracellular Traps Is Not a Useful Biomarker for Assessing Disease Activity in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. [PLoS One. 11 \(2\): e0148197.](#)
10. Rai, A.K. *et al.* (2017) Exonal switch down-regulates the expression of CD5 on blasts of acute T cell leukaemia. [Clin Exp Immunol. 190 \(3\): 340-350.](#)
11. Hoppenbrouwers, T. *et al.* (2018) Neutrophil Extracellular Traps in Children With Meningococcal Sepsis. [Pediatr Crit Care Med. 19 \(6\): e286-e291.](#)
12. Helseth, R. *et al.* (2019) Glucose associated NETosis in patients with ST-elevation myocardial infarction: an observational study. [BMC Cardiovasc Disord. 19 \(1\): 221.](#)
13. Manfredi, A.A. *et al.* (2021) Platelet phagocytosis via PSGL1 and accumulation of microparticles in systemic sclerosis. [Arthritis Rheumatol. Jul 19 \[Epub ahead of print\].](#)
14. de Moraes Mazetto, B. *et al.* (2022) Association between neutrophil extracellular traps (NETs) and thrombosis in antiphospholipid syndrome [Thrombosis Res. 214: 132-7.](#)
15. Robinson, H. *et al.* (2022) The effect of expressive writing on wound healing: Immunohistochemistry analysis of skin tissue two weeks after punch biopsy wounding. [J Psychosom Res. 161: 110987.](#)
16. Zhu, D. *et al.* (2024) Neutrophil activation biomarker pentraxin 3 for diagnosis and monitoring of macrophage activation syndrome occurrence in adult-onset Still's disease. [J Autoimmun. 144: 103182.](#)
17. Rogato, F. *et al.* (2024) Leukemia cutis as a prominent clinical sign in a dog with acute myeloid leukemia. [Vet Clin Pathol. 53 \(4\): 448-57.](#)

---

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1757T>

**Regulatory**

For research purposes only

**Related Products****Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>

**Recommended Negative Controls**[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M437190:250310'

**Printed on 27 Mar 2025**