Datasheet: MCA1765GT BATCH NUMBER 0412

Description:	MOUSE ANTI HUMAN CD70
Specificity:	CD70
Other names:	TNFSF7
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	Bu69
Isotype:	lgG1
Quantity:	25 µg

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>rad-antibodies.com/protocols.

		0013.			
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry				1/50 - 1/200
	Immunohistology - Frozen				
	Immunohistology - Paraffin			•	
	ELISA			•	
	Immunoprecipitation			•	
	Western Blotting			•	
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.				
Target Species	Human				
Product Form	Purified IgG - liquid				
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant			m tissue culture	
Buffer Solution	Phosphate buffered salin	e			
Preservative Stabilisers	0.09% Sodium Azide				

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
External Database Links	UniProt: P32970 Related reagents
	Entrez Gene: <u>970</u> CD70 <u>Related reagents</u>
Synonyms	CD27L, CD27LG, TNFSF7
RRID	AB_2075213
Specificity	Mouse anti Human CD70 antibody, clone Bu69 recognises human CD70, also known as CD27 ligand (CD27-L) or Tumor necrosis factor ligand superfamily member 7. CD70 is a 193 amino acid ~29 kDa single pass type II transmembrane glycoprotein belonging to the tumour necrosis factor (TNF) family (<u>UniProt: P32970</u>).
	CD70 is expressed on a small number of activated T-cells and the majority of activated B-cells (<u>Bowman <i>et al.</i> 1994</u>). Interaction of CD70 with its receptor, CD27, is thought to play an important role in lymphocyte activation, proliferation, survival and differentiation (<u>Hintzen <i>et al.</i> 1994</u>).
	Mouse anti Human CD70 antibody, clone Bu69 has been used for the inhibition of neural stem cell induced apoptosis of allogeneic T cells <i>in vitro</i> (<u>Lee <i>et al.</i> 2013</u>).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Garland Publishing. (1997) Leucocyte typing VI. White cell differentiation antigens. p522 - 4
	 Zambello, R. <i>et al.</i> (2000) Analysis of TNF-receptor and ligand superfamily molecules in patients with lymphoproliferative disease of granular lymphocytes. <u>Blood. 96 (2): 647-54.</u> Lee, E.M. <i>et al.</i> (2013) CD70-CD27 ligation between neural stem cells and CD4+ T cells induces Fas-FasL-mediated T-cell death. <u>Stem Cell Res Ther. 4 (3): 56.</u>
Storage	Store at +4°C or at -20°C if preferred.
	This product should be stored undiluted.
	Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	18 months from date of despatch.
Health And Safety	Material Safety Datasheet documentation #10040 available at:

Information	https://www.bio-rad-antibodies.com/SDS/MCA1765GT 10040		
Regulatory	For research purposes only		

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12)	RPE			
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>				
Goat Anti Mouse IgG (STAR76)	RPE			
Rabbit Anti Mouse IgG (STAR13)	HRP			
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>			
Goat Anti Mouse IgG (H/L) (STAR117)	<u>Alk. Phos., DyLight®488, DyLight®550,</u>			
	<u>DyLight®650</u> , <u>DyLight®680</u> , <u>DyLight®800</u> ,			
	<u>FITC, HRP</u>			
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>			
Goat Anti Mouse IgG (STAR77)	HRP			
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP			
Recommended Negative Controls				

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M343003:190110'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint