

Datasheet: MCA1767

BATCH NUMBER 158692

Description:	RAT ANTI MOUSE CD4
Specificity:	CD4
Other names:	L3T4 ANTIGEN, LY-4
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	YTS191.1
Isotype:	IgG2b
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	<p>UniProt: P06332 Related reagents</p> <p>Entrez Gene: 12504 Cd4 Related reagents</p>
RRID	AB_322769
Specificity	<p>Rat anti Mouse CD4 antibody, clone YTS191.1 recognizes the murine CD4 cell surface antigen, expressed by a subset of T lymphocytes.</p> <p>Rat anti Mouse CD4 antibody, clone YTS191.1 exhibits depleting activity when used <i>in vivo</i> (Bemelman <i>et al.</i> 1998).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Cobbold, S.P. <i>et al.</i> (1990) The induction of skin graft tolerance in major histocompatibility complex-mismatched or primed recipients: primed T cells can be tolerized in the periphery with anti-CD4 and anti-CD8 antibodies. Eur J Immunol. 20 (12): 2747-55. 2. Bemelman, F. <i>et al.</i> (1998) Bone marrow transplantation induces either clonal deletion or infectious tolerance depending on the dose. J Immunol. 160 (6): 2645-8. 3. Higgins, L.M. <i>et al.</i> (1999) Regulation of T cell activation in vitro and in vivo by targeting the OX40-OX40 ligand interaction: amelioration of ongoing inflammatory bowel disease with an OX40-IgG fusion protein, but not with an OX40 ligand-IgG fusion protein. J Immunol. 162 (1): 486-93. 4. Croxford, J.L. <i>et al.</i> (2001) Different therapeutic outcomes in experimental allergic encephalomyelitis dependent upon the mode of delivery of IL-10: a comparison of the effects of protein, adenoviral or retroviral IL-10 delivery into the central nervous system. J Immunol. 166: 4124-30. 5. Eller, K. <i>et al.</i> (2011) IL-9 production by regulatory T cells recruits mast cells that are essential for regulatory T cell-induced immune suppression. J Immunol. 186: 83-91. 6. Gaupp, S. <i>et al.</i> (2008) Amelioration of experimental autoimmune encephalomyelitis in IL-4Ralpha^{-/-} mice implicates compensatory up-regulation of Th2-type cytokines. Am J Pathol. 173: 119-29. 7. Grimm, M. <i>et al.</i> (2010) Evaluation of immunological escape mechanisms in a mouse model of colorectal liver metastases. BMC Cancer. 10: 82. 8. Jégou, J.F. <i>et al.</i> (2007) C3d Binding to the Myelin Oligodendrocyte Glycoprotein

Results in an Exacerbated Experimental Autoimmune Encephalomyelitis [J Immunol. 178: 3323-31.](#)

9. Huber, J.M. *et al.* (2009) The proteasome inhibitor bortezomib aggravates renal ischemia-reperfusion injury. [Am J Physiol Renal Physiol. 297: F451-60.](#)

10. Wolf, D. *et al.* (2005) CD4+CD25+ regulatory T cells inhibit experimental anti-glomerular basement membrane glomerulonephritis in mice. [J Am Soc Nephrol. 16: 1360-70.](#)

11. Abdulreda, M.H. *et al.* (2011) High-resolution, noninvasive longitudinal live imaging of immune responses. [Proc Natl Acad Sci U S A. 108: 12863-8.](#)

12. Nakashima, H. *et al.* (2011) A Novel Combination Immunotherapy for Cancer by IL-13R α 2-Targeted DNA Vaccine and Immunotoxin in Murine Tumor Models. [J Immunol. 187: 4935-46.](#)

13. Scotland, R.S. *et al.* (2011) Sex-differences in resident immune cell phenotype underlies more efficient acute inflammatory responses in female mice. [Blood. 118: 5918-27.](#)

14. Zitt, E. *et al.* (2011) The selective mineralocorticoid receptor antagonist eplerenone is protective in mild anti-GBM glomerulonephritis. [Int J Clin Exp Pathol. 4:606-15.](#)

15. Nelvagal, H.R. *et al.* (2020) Comparative proteomic profiling reveals mechanisms for early spinal cord vulnerability in CLN1 disease. [Sci Rep. 10 \(1\): 15157.](#)

16. Groh, J. *et al.* (2021) Immune modulation attenuates infantile neuronal ceroid lipofuscinosis in mice before and after disease onset [Brain Communications. fcab047.](#)

17. Karikari, A.A. *et al.* (2022) Neurodegeneration by α -synuclein-specific T cells in AAV-A53T- α -synuclein Parkinson's disease mice. [Brain Behav Immun. 101: 194-210.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1767>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69...)

[FITC](#)

Goat Anti Rat IgG (STAR73...)

[RPE](#)

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) [DyLight@550](#), [DyLight@650](#), [DyLight@800](#)

Rabbit Anti Rat IgG (STAR21...)

[HRP](#)

Rabbit Anti Rat IgG (STAR16...)

[DyLight@800](#)

Goat Anti Rat IgG (STAR131...)

[Alk. Phos.](#), [Biotin](#)

Rabbit Anti Rat IgG (STAR17...)

[FITC](#)

Goat Anti Rat IgG (STAR72...)

[HRP](#)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M383154:210513'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)