

## Datasheet: MCA1768A405

|                      |                                     |
|----------------------|-------------------------------------|
| <b>Description:</b>  | RAT ANTI MOUSE CD8:Alexa Fluor® 405 |
| <b>Specificity:</b>  | CD8                                 |
| <b>Format:</b>       | ALEXA FLUOR® 405                    |
| <b>Product Type:</b> | Monoclonal Antibody                 |
| <b>Clone:</b>        | YTS169.4                            |
| <b>Isotype:</b>      | IgG2b                               |
| <b>Quantity:</b>     | 100 TESTS/1ml                       |

## Product Details

**RRID** AB\_566911

**Applications** This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

|                | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ■   |    |                | 1/5 - 1/10         |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls

**Target Species** Mouse

**Product Form** Purified IgG conjugated to Alexa Fluor® 405 - liquid

| Max Ex/Em | Fluorophore     | Excitation Max (nm) | Emission Max (nm) |
|-----------|-----------------|---------------------|-------------------|
|           | Alexa Fluor®405 | 401                 | 421               |

**Preparation** Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

**Buffer Solution** Phosphate buffered saline

**Preservative** 0.09% Sodium Azide  
**Stabilisers** 1% Bovine Serum Albumin

**Approx. Protein Concentrations** IgG concentration 0.05 mg/ml

**External Database Links**

**UniProt:**  
[P01731](#) [Related reagents](#)  
[P10300](#) [Related reagents](#)

**Entrez Gene:**

[12525](#) Cd8a [Related reagents](#)

[12526](#) Cd8b1 [Related reagents](#)

---

**Synonyms**

Cd8b1, Ly-3, Lyt2, Lyt-2, Lyt3, Lyt-3

---

**Specificity**

**Rat anti Mouse CD8 antibody, clone YTS169.4** recognizes the murine CD8 cell surface antigen, expressed by a subset of T lymphocytes.

Rat anti Mouse CD8 antibody, clone YTS169.4 exhibits depleting activity when used *in vivo*.

---

**Flow Cytometry**

Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#))

---

**References**

1. Cobbold, S.P. *et al.* (1990) The induction of skin graft tolerance in major histocompatibility complex-mismatched or primed recipients: primed T cells can be tolerized in the periphery with anti-CD4 and anti-CD8 antibodies. [Eur J Immunol. 20 \(12\): 2747-55.](#)
2. Bemelman, F. *et al.* (1998) Bone marrow transplantation induces either clonal deletion or infectious tolerance depending on the dose. [J Immunol. 160 \(6\): 2645-8.](#)
3. Cobbold SP *et al.* (1984) Therapy with monoclonal antibodies by elimination of T-cell subsets *in vivo*. [Nature. 312 \(5994\): 548-51.](#)
4. Wise, M.P. *et al.* (1998) Linked suppression of skin graft rejection can operate through indirect recognition. [J Immunol. 161 \(11\): 5813-6.](#)
5. Higgins, L.M. *et al.* (1999) Regulation of T cell activation in vitro and in vivo by targeting the OX40-OX40 ligand interaction: amelioration of ongoing inflammatory bowel disease with an OX40-IgG fusion protein, but not with an OX40 ligand-IgG fusion protein. [J Immunol. 162 \(1\): 486-93.](#)
6. Scotland, R.S. *et al.* (2011) Sex-differences in resident immune cell phenotype underlies more efficient acute inflammatory responses in female mice. [Blood. 118: 5918-27.](#)
7. Matsubara, K. *et al.* (2016) Immune activation during the implantation phase causes preeclampsia-like symptoms via the CD40-CD40 ligand pathway in pregnant mice. [Hypertens Res. 39 \(6\): 407-14.](#)
8. Jaffar, Z. *et al.* (2002) A key role for prostaglandin I2 in limiting lung mucosal Th2, but not Th1, responses to inhaled allergen. [J Immunol. 169 \(10\): 5997-6004.](#)
9. Zirger, J.M. *et al.* (2012) Immune-mediated loss of transgene expression from virally transduced brain cells is irreversible, mediated by IFN $\gamma$ , perforin, and TNF $\alpha$ , and due to the elimination of transduced cells. [Mol Ther. 20 \(4\): 808-19.](#)
10. Abd-elhakim, Y.M. *et al.* (2016) Hemato-immunologic impact of subchronic exposure to melamine and/or formaldehyde in mice. [J Immunotoxicol. 13 \(5\): 713-22.](#)
11. Ismail, S.A.A. (2017) Ameliorative Potential of *Spirulina platensis* against Lead Acetate Induced Immuno-Suppression and Kidney Apoptosis in Rats [Ann Clin Pathol 5\(5\): 1120.](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use

---

**Shelf Life** 18 months from date of despatch

---

**Acknowledgements** This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or [outlicensing@thermofisher.com](mailto:outlicensing@thermofisher.com)

---

**Health And Safety Information** Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M341594:190109'

**Printed on 11 Jan 2019**

---

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)