

Datasheet: MCA1768PET

Description:	RAT ANTI MOUSE CD8:RPE
Specificity:	CD8
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	YTS169.4
Isotype:	IgG2b
Quantity:	25 TESTS/0.25ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse						
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>RPE 488nm laser</td> <td>496</td> <td>578</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	RPE 488nm laser	496	578
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
RPE 488nm laser	496	578					
Preparation	Purified IgG prepared by affinity chromatography on Protein G						
Buffer Solution	Phosphate buffered saline						
Preservative	0.09% Sodium Azide						
Stabilisers	1% Bovine Serum Albumin 5% Sucrose						

External Database Links

UniProt:

[P01731](#) [Related reagents](#)
[P10300](#) [Related reagents](#)

Entrez Gene:

[12525](#) Cd8a [Related reagents](#)
[12526](#) Cd8b1 [Related reagents](#)

Synonyms	Cd8b1, Ly-3, Lyt2, Lyt-2, Lyt3, Lyt-3
Specificity	<p>Rat anti Mouse CD8 antibody, clone YTS169.4 recognizes the murine CD8 cell surface antigen, expressed by a subset of T lymphocytes.</p> <p>Rat anti Mouse CD8 antibody, clone YTS169.4 exhibits depleting activity when used <i>in vivo</i>.</p>
Flow Cytometry	<p>Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.</p> <p>The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).</p>
References	<ol style="list-style-type: none"> Cobbold, S.P. <i>et al.</i> (1990) The induction of skin graft tolerance in major histocompatibility complex-mismatched or primed recipients: primed T cells can be tolerized in the periphery with anti-CD4 and anti-CD8 antibodies. Eur J Immunol. 20 (12): 2747-55. Bemelman, F. <i>et al.</i> (1998) Bone marrow transplantation induces either clonal deletion or infectious tolerance depending on the dose. J Immunol. 160 (6): 2645-8. Cobbold SP <i>et al.</i> (1984) Therapy with monoclonal antibodies by elimination of T-cell subsets <i>in vivo</i>. Nature. 312 (5994): 548-51. Wise, M.P. <i>et al.</i> (1998) Linked suppression of skin graft rejection can operate through indirect recognition. J Immunol. 161 (11): 5813-6. Higgins, L.M. <i>et al.</i> (1999) Regulation of T cell activation in vitro and in vivo by targeting the OX40-OX40 ligand interaction: amelioration of ongoing inflammatory bowel disease with an OX40-IgG fusion protein, but not with an OX40 ligand-IgG fusion protein. J Immunol. 162 (1): 486-93. Scotland, R.S. <i>et al.</i> (2011) Sex-differences in resident immune cell phenotype underlies more efficient acute inflammatory responses in female mice. Blood. 118: 5918-27. Matsubara, K. <i>et al.</i> (2016) Immune activation during the implantation phase causes preeclampsia-like symptoms via the CD40-CD40 ligand pathway in pregnant mice. Hypertens Res. 39 (6): 407-14. Jaffar, Z. <i>et al.</i> (2002) A key role for prostaglandin I2 in limiting lung mucosal Th2, but not Th1, responses to inhaled allergen. J Immunol. 169 (10): 5997-6004. Zirger, J.M. <i>et al.</i> (2012) Immune-mediated loss of transgene expression from virally transduced brain cells is irreversible, mediated by IFNγ, perforin, and TNFα, and due to the elimination of transduced cells. Mol Ther. 20 (4): 808-19. Abd-elhakim, Y.M. <i>et al.</i> (2016) Hemato-immunologic impact of subchronic exposure to melamine and/or formaldehyde in mice. J Immunotoxicol. 13 (5): 713-22. Ismail, S.A.A. (2017) Ameliorative Potential of <i>Spirulina platensis</i> against Lead Acetate Induced Immuno-Suppression and Kidney Apoptosis in Rats Ann Clin Pathol 5(5): 1120.
Storage	<p>Store at +4°C.</p> <p>DO NOT FREEZE.</p> <p>This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Shelf Life	12 months from date of despatch.
Health And Safety Information	<p>Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf</p>

**North & South
America**

Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M331635:180924'

Printed on 06 Dec 2018
