

Datasheet: MCA1768T

| | |
|----------------------|---------------------|
| Description: | RAT ANTI MOUSE CD8 |
| Specificity: | CD8 |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | YTS169.4 |
| Isotype: | IgG2b |
| Quantity: | 25 µg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | 1/50 - 1/100 |
| Immunohistology - Frozen | ▪ | | | |
| Immunohistology - Paraffin | | | ▪ | |
| ELISA | | | ▪ | |
| Immunoprecipitation | | | ▪ | |
| Western Blotting | | | ▪ | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| | |
|---------------------------------|---|
| Target Species | Mouse |
| Product Form | Purified IgG - liquid |
| Preparation | Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant |
| Buffer Solution | Phosphate buffered saline |
| Preservative Stabilisers | 0.09% Sodium Azide |
| Carrier Free | Yes |

| | |
|---------------------------------------|---|
| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml |
| External Database Links | <p>UniProt:</p> <p>P01731 Related reagents</p> <p>P10300 Related reagents</p> <p>Entrez Gene:</p> <p>12525 Cd8a Related reagents</p> <p>12526 Cd8b1 Related reagents</p> |
| Synonyms | Cd8b1, Ly-3, Lyt2, Lyt-2, Lyt3, Lyt-3 |
| RRID | AB_1102357 |
| Specificity | <p>Rat anti Mouse CD8 antibody, clone YTS169.4 recognizes the murine CD8 cell surface antigen, expressed by a subset of T lymphocytes.</p> <p>Rat anti Mouse CD8 antibody, clone YTS169.4 exhibits depleting activity when used <i>in vivo</i>.</p> |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. |
| References | <ol style="list-style-type: none"> 1. Cobbold, S.P. <i>et al.</i> (1990) The induction of skin graft tolerance in major histocompatibility complex-mismatched or primed recipients: primed T cells can be tolerized in the periphery with anti-CD4 and anti-CD8 antibodies. Eur J Immunol. 20 (12): 2747-55. 2. Bemelman, F. <i>et al.</i> (1998) Bone marrow transplantation induces either clonal deletion or infectious tolerance depending on the dose. J Immunol. 160 (6): 2645-8. 3. Cobbold SP <i>et al.</i> (1984) Therapy with monoclonal antibodies by elimination of T-cell subsets <i>in vivo</i>. Nature. 312 (5994): 548-51. 4. Wise, M.P. <i>et al.</i> (1998) Linked suppression of skin graft rejection can operate through indirect recognition. J Immunol. 161 (11): 5813-6. 5. Higgins, L.M. <i>et al.</i> (1999) Regulation of T cell activation in vitro and in vivo by targeting the OX40-OX40 ligand interaction: amelioration of ongoing inflammatory bowel disease with an OX40-IgG fusion protein, but not with an OX40 ligand-IgG fusion protein. J Immunol. 162 (1): 486-93. 6. Scotland, R.S. <i>et al.</i> (2011) Sex-differences in resident immune cell phenotype underlies more efficient acute inflammatory responses in female mice. Blood. 118: 5918-27. 7. Matsubara, K. <i>et al.</i> (2016) Immune activation during the implantation phase causes preeclampsia-like symptoms via the CD40-CD40 ligand pathway in pregnant mice. Hypertens Res. 39 (6): 407-14. 8. Jaffar, Z. <i>et al.</i> (2002) A key role for prostaglandin I2 in limiting lung mucosal Th2, but not Th1, responses to inhaled allergen. J Immunol. 169 (10): 5997-6004. 9. Zirger, J.M. <i>et al.</i> (2012) Immune-mediated loss of transgene expression from virally transduced brain cells is irreversible, mediated by IFNγ, perforin, and TNFα, and due to |

the elimination of transduced cells. [Mol Ther. 20 \(4\): 808-19.](#)

10. Abd-elhakim, Y.M. *et al.* (2016) Hemato-immunologic impact of subchronic exposure to melamine and/or formaldehyde in mice. [J Immunotoxicol. 13 \(5\): 713-22.](#)

11. Lejeune, P. *et al.* (2021) Immunostimulatory effects of targeted thorium-227 conjugates as single agent and in combination with anti-PD-L1 therapy. [J Immunother Cancer. \(10\):e002387.](#)

12. Nelvagal, H.R. *et al.* (2020) Comparative proteomic profiling reveals mechanisms for early spinal cord vulnerability in CLN1 disease. [Sci Rep. 10 \(1\): 15157.](#)

13. de Souza, T.A. *et al.* (2018) Relationship between the inflammatory tumor microenvironment and different histologic types of canine mammary tumors. [Res Vet Sci. 119: 209-14.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

| | |
|--|---|
| Goat Anti Rat IgG (STAR69...) | FITC |
| Goat Anti Rat IgG (STAR73...) | RPE |
| Rabbit Anti Rat IgG (STAR17...) | FITC |
| Goat Anti Rat IgG (STAR131...) | Alk. Phos. , Biotin |
| Goat Anti Rat IgG (STAR72...) | HRP |
| Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) | DyLight@550 , DyLight@650 , DyLight@800 |
| Rabbit Anti Rat IgG (STAR21...) | HRP |
| Rabbit Anti Rat IgG (STAR16...) | DyLight@800 |

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M383082:210513'

Printed on 16 Sep 2022