

Datasheet: MCA1774F

Description:	MOUSE ANTI DOG CD3:FITC
Specificity:	CD3
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	CA17.2A12
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

RRID AB_2291174

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Dog

Product Form Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Preparation Purified IgG prepared by affinity chromatography on Protein G

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide
Stabilisers 1% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.1 mg/ml

Immunogen Affinity enriched TCR/CD3 membrane proteins isolated from thymocytes and the T cell line CLGL-90

External Database Links **UniProt:**

Entrez Gene:

[442981](#) CD3E [Related reagents](#)

Specificity

Mouse anti Dog CD3 antibody, clone CA17.2A12 recognizes the canine CD3 cell surface antigen, expressed by thymocytes and mature T lymphocytes. CD3 is engaged in the surface expression of the T-cell antigen receptor (TCR) and the signal transduction pathway resulting from MHC ligand binding to the TCR. CD3 is made up of a number of invariant subchains of the immunoglobulin superfamily.

Mouse anti Dog CD3 clone CA17.2A12 is a valuable flow cytometric and immunohistologic tool for canine lymphoma detection of T-cell origin ([Miniscalco et al. 2003](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells or 100ul whole blood.

N.B. MCA1774F should not be used with MCA1781PE (mouse anti canine B-cells), in dual colour flow cytometry, due to non-specific interactions between the two reagents.

References

1. Moore, P.F. and Rossitto, P.V. (1993) Development of monoclonal antibodies to canine T cell receptor complex (TCR/CD3) and their utilisation in the diagnosis of T cell neoplasia. [Vet. Pathol. 30: 457. Abstract 117](#)
2. McDonough, S. P. and Moore, P. F. (2000) Clinical, hematologic, and immunophenotypic characterization of canine large granular lymphocytosis. [Vet Pathol. 37:637-46.](#)
3. Moore, P.F. et al. (2006) Canine hemophagocytic histiocytic sarcoma: a proliferative disorder of CD11d+ macrophages. [Vet Pathol. 43 \(5\): 632-45.](#)
4. Vernau, W and Moore, P. F. (1999) An immunophenotypic study of canine leukemias and preliminary assessment of clonality by polymerase chain reaction. [Vet Immunol Immunopathol. 69:145-64.](#)
5. Moreno, J. et al (1999) The immune response and PBMC subsets in canine visceral leishmaniasis before, and after, chemotherapy. [Vet Immunol Immunopathol. 71:181-95.](#)
6. Machado, G.F. et al. (2011) Intravascular Lymphomatosis in the Central Nervous System of Dogs: Immunohistochemical Investigation in Two Cases [Braz J Vet Pathol 4: 47-51](#)
7. Fellman, C.L. et al. (2011) Cyclosporine A affects the *in vitro* expression of T cell activation-related molecules and cytokines in dogs. [Vet Immunol Immunopathol. 140: 175-80.](#)
8. Watabe, A. et al. (2011) Alterations of lymphocyte subpopulations in healthy dogs with aging and in dogs with cancer. [Vet Immunol Immunopathol. 142: 189-200.](#)
9. Hsiao, Y.W. et al (2004) Tumor-infiltrating lymphocyte secretion of IL-6 antagonizes tumor-derived TGF-beta 1 and restores the lymphokine-activated killing activity. [J Immunol. 172: 1508-14.](#)
10. Huang, Y.C. et al. (2008) CD5-low expression lymphocytes in canine peripheral blood show characteristics of natural killer cells. [J Leukoc Biol. 84: 1501-10.](#)
11. Out, T.A. et al. (2002) Local T-cell activation after segmental allergen challenge in the lungs of allergic dogs. [Immunology. 105: 499-508.](#)
12. Zentek, J. et al. (2002) Morphology and immunopathology of the small and large intestine in dogs with nonspecific dietary sensitivity. [J Nutr. 132: 1652S-4S.](#)
13. Hai, M. et al. (2008) Potential genotoxicity from integration sites in CLAD dogs treated successfully with gammaretroviral vector-mediated gene therapy. [Gene Ther. 15: 1067-71.](#)
14. Altmann, S. et al. (2008) High Mobility Group Box 1-Protein expression in canine haematopoietic cells and influence on canine peripheral blood mononuclear cell proliferative activity [Vet Immunol Immunopathol. 126: 367-72.](#)
15. Ting-De Ravin, S.S. et al. (2006) Correction of canine X-linked severe combined immunodeficiency by in vivo retroviral gene therapy. [Blood. 107: 3091-7.](#)

16. Miranda, S. *et al.* (2007) Characterization of circulating lymphocyte subpopulations in canine leishmaniasis throughout treatment with antimonials and allopurinol. [Vet Parasitol. 144: 251-60.](#)
17. Maiolini, A. *et al.* (2012) Toll-like receptors 4 and 9 are responsible for the maintenance of the inflammatory reaction in canine steroid-responsive meningitis-arteritis, a large animal model for neutrophilic meningitis. [J Neuroinflammation. 9: 226.](#)
18. Aresu, L. *et al.* (2014) VEGF and MMP-9: biomarkers for canine lymphoma. [Vet Comp Oncol. 12: 29-36.](#)
19. Schaut, R.G. *et al.* (2016) Regulatory IgDhi B Cells Suppress T Cell Function via IL-10 and PD-L1 during Progressive Visceral Leishmaniasis. [J Immunol. 196 \(10\): 4100-9.](#)
20. Villaescusa A *et al.* (2012) Evaluation of peripheral blood lymphocyte subsets in family-owned dogs naturally infected by *Ehrlichia canis*. [Comp Immunol Microbiol Infect Dis. 35 \(4\): 391-6.](#)
21. Riondato, F. *et al.* (2016) Analytical and diagnostic validation of a flow cytometric strategy to quantify blood and marrow infiltration in dogs with large B-cell lymphoma. [Cytometry B Clin Cytom. 90 \(6\): 525-530.](#)
22. Byrne, K. *et al.* (2000) A standardized gating technique for the generation of flow cytometry data for normal canine and normal feline blood lymphocytes. [Vet Immunol Immunopathol. 73:167-82.](#)
23. Perosso, J. *et al.* (2014) Alteration of sFAS and sFAS ligand expression during canine visceral leishmaniasis. [Vet Parasitol. 205 \(3-4\): 417-23.](#)
24. Grøndahl-Rosado C *et al.* (2015) NCR1+ cells in dogs show phenotypic characteristics of natural killer cells. [Vet Res Commun. 39 \(1\): 19-30.](#)
25. Miller, J. *et al.* (2015) Humoral and Cellular Immune Response in Canine Hypothyroidism. [J Comp Pathol. 153 \(1\): 28-37.](#)
26. McGill, J.L. *et al.* (2016) Vaccination with an Attenuated Mutant of *Ehrlichia chaffeensis* Induces Pathogen-Specific CD4+ T Cell Immunity and Protection from Tick-Transmitted Wild-Type Challenge in the Canine Host. [PLoS One. 11 \(2\): e0148229.](#)
27. Constantinoiu CC *et al.* (2015) Mucosal tolerance of the hookworm *Ancylostoma caninum* in the gut of naturally infected wild dogs. [Parasite Immunol. Jul 27 \[Epub ahead of print\]](#)
28. Duz, A.L. *et al.* (2014) The TcI and TcII *Trypanosoma cruzi* experimental infections induce distinct immune responses and cardiac fibrosis in dogs. [Mem Inst Oswaldo Cruz. 109 \(8\): 1005-13.](#)
29. Mie, K. *et al.* (2016) Change in peripheral blood lymphocyte count in dogs following adoptive immunotherapy using lymphokine-activated T killer cells combined with palliative tumor resection. [Vet Immunol Immunopathol. 177: 58-63.](#)
30. Schaut, R.G. *et al.* (2016) Recovery of antigen-specific T cell responses from dogs infected with *Leishmania (L.) infantum* by use of vaccine associated TLR-agonist adjuvant. [Vaccine. 34 \(44\): 5225-34.](#)
31. Gelain, M.E. *et al.* (2014) CD44 in canine leukemia: analysis of mRNA and protein expression in peripheral blood. [Vet Immunol Immunopathol. 159 \(1-2\): 91-6.](#)
32. Michael, H.T. *et al.* (2013) Isolation and characterization of canine natural killer cells. [Vet Immunol Immunopathol. 155 \(3\): 211-7.](#)
33. Bonnefont-Rebeix, C. *et al.* (2016) Characterization of a novel canine T-cell line established from a spontaneously occurring aggressive T-cell lymphoma with large granular cell morphology. [Immunobiology. 221 \(1\): 12-22.](#)
34. Schmidli, M.R. *et al.* (2018) Inflammatory pattern of the infrapatellar fat pad in dogs with canine cruciate ligament disease. [BMC Vet Res. 14 \(1\): 161.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product

contain a precipitate we recommend microcentrifugation before use.

Guarantee	18 months from date of despatch.
------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M334453:181203'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)