



Datasheet: MCA1778S

BATCH NUMBER 150750

Description:	MOUSE ANTI DOG CD11c
Specificity:	CD11c
Other names:	INTEGRIN ALPHA X CHAIN
Format:	S/N
Product Type:	Monoclonal Antibody
Clone:	CA11.6A1
Isotype:	IgG1
Quantity:	2 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species

Dog

Species Cross Reactivity

Reacts with: Hooded Seal, Raccoon

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form	Tissue Culture Supernatant - liquid
Preservative Stabilisers	0.1% Sodium Azide
RRID	AB_322942
Specificity	<p>Mouse anti Dog CD11c antibody, clone CA11.6A1 recognizes the canine CD11c cell surface antigen, a member of the alpha integrin family. Canine CD11c is expressed by monocytes, granulocytes and by dendritic cells.</p> <p>Mouse anti Dog CD11c, clone CA11.6A1 immunoprecipitates proteins of approximately 95 kDa, corresponding to the common β chain of the CD11/CD18 heterodimer and ~150 kDa, the CD11c; chain from canine leukocyte preparations (Danilenko et al. 1992)</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10^6 cells or 100ul whole blood
References	<ol style="list-style-type: none"> Danilenko, D.M. <i>et al.</i> (1992) Canine leukocyte cell adhesion molecules (LeuCAMs): characterization of the CD11/CD18 family. Tissue Antigens 40: 13-21. Kang, J.W. <i>et al.</i> (2008) Soluble factors-mediated immunomodulatory effects of canine adipose tissue-derived mesenchymal stem cells. Stem Cells Dev. 17: 681-93. Affolter, V.K. and Moore, P.F. (2002) Localized and disseminated histiocytic sarcoma of dendritic cell origin in dogs. Vet Pathol. 39: 74-83. Bird, R.C. <i>et al.</i> (2008) An allogeneic hybrid-cell fusion vaccine against canine mammary cancer. Vet Immunol Immunopathol. 123: 289-304. Catchpole, B. <i>et al.</i> (2002) Generation of blood-derived dendritic cells in dogs with oral malignant melanoma. J Comp Pathol. 126: 238-41. Isotani, M. <i>et al.</i> (2006) Efficient generation of canine bone marrow-derived dendritic cells. J Vet Med Sci. 68: 809-14. Liu, C.C. <i>et al.</i> (2008) Transient downregulation of monocyte-derived dendritic-cell differentiation, function, and survival during tumoral progression and regression in an in vivo canine model of transmissible venereal tumor. Cancer Immunol Immunother. 57: 479-91. McDonough, S.P. and Moore, P.F. (2000) Clinical, hematologic, and immunophenotypic characterization of canine large granular lymphocytosis. Vet Pathol. 37: 637-46. Wang, Y.S. <i>et al.</i> (2007) Characterization of canine monocyte-derived dendritic cells with phenotypic and functional differentiation. Can J Vet Res. 71: 165-74. Mathes, M. <i>et al.</i> (2006) Evaluation of liposomal clodronate in experimental spontaneous autoimmune hemolytic anemia in dogs. Exp Hematol. 34: 1393-402. Sanchez, M.A. <i>et al.</i> (2004) Organ-specific immunity in canine visceral leishmaniasis: analysis of symptomatic and asymptomatic dogs naturally infected with <i>Leishmania chagasi</i>. Am J Trop Med Hyg. 70: 618-24. Ricklin Gutzwiller, M.E. <i>et al.</i> (2010) Comparative analysis of canine monocyte- and bone-marrow-derived dendritic cells. Vet Res. 41: 40. Ibisch, C. <i>et al.</i> (2005) Functional canine dendritic cells can be generated in vitro from peripheral blood mononuclear cells and contain a cytoplasmic ultrastructural marker. J Immunol Methods. 298: 175-82. Wang, Y.S. <i>et al.</i> (2008) Cytokine profiles of canine monocyte-derived dendritic cells

as a function of lipopolysaccharide- or tumor necrosis factor-alpha-induced maturation. [Vet Immunol Immunopathol. 118: 186-98.](#)

15. Schwartz, M. *et al.* (2008) Selective CD11a upregulation on neutrophils in the acute phase of steroid-responsive meningitis-arteritis in dogs. [Vet Immunol Immunopathol. 126: 248-55.](#)

16. Pai, C.C. *et al.* (2011) Immunopathogenic behaviors of canine transmissible venereal tumor in dogs following an immunotherapy using dendritic/tumor cell hybrid. [Vet Immunol Immunopathol. 139 \(2-4\): 187-99.](#)

17. Figueiredo, M.M. *et al.* (2013) Expression of Toll-like Receptors 2 and 9 in cells of dog jejunum and colon naturally infected with *Leishmania infantum*. [BMC Immunol. 14: 22.](#)

18. Larsen, A.K. *et al.* (2013) Entry and elimination of marine mammal *Brucella* spp. by hooded seal (*Cystophora cristata*) alveolar macrophages *in vitro*. [PLoS One. 8: e70186.](#)

19. Heinrich, F. *et al.* (2015) Immunophenotyping of immune cell populations in the raccoon (*Procyon lotor*). [Vet Immunol Immunopathol. 168 \(3-4\): 140-6.](#)

20. Paoloni, M. *et al.* (2015) Defining the Pharmacodynamic Profile and Therapeutic Index of NHS-IL12 Immunocytokine in Dogs with Malignant Melanoma. [PLoS One. 10 \(6\): e0129954.](#)

21. Bonnefont-Rebeix, C. *et al.* (2016) Characterization of a novel canine T-cell line established from a spontaneously occurring aggressive T-cell lymphoma with large granular cell morphology. [Immunobiology. 221 \(1\): 12-22.](#)

22. Constantinoiu, C.C. *et al.* (2015) Mucosal tolerance of the hookworm *Ancylostoma caninum* in the gut of naturally infected wild dogs. [Parasite Immunol. Jul 27 \[Epub ahead of print\].](#)

23. Stokol, T. *et al.* (2015) Alkaline phosphatase is a useful cytochemical marker for the diagnosis of acute myelomonocytic and monocytic leukemia in the dog. [Vet Clin Pathol. 44 \(1\): 79-93.](#)

24. Heinrich, F. *et al.* (2015) Passage-dependent morphological and phenotypical changes of a canine histiocytic sarcoma cell line (DH82 cells). [Vet Immunol Immunopathol. 163 \(1-2\): 86-92.](#)

25. Qeska, V. *et al.* (2014) Canine distemper virus infection leads to an inhibitory phenotype of monocyte-derived dendritic cells *in vitro* with reduced expression of co-stimulatory molecules and increased interleukin-10 transcription. [PLoS One. 9 \(4\): e96121.](#)

26. Bird, R.C. *et al.* (2019) Autologous hybrid cell fusion vaccine in a spontaneous intermediate model of breast carcinoma. [J Vet Sci. 20 \(5\): e48.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10336 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1778S>
10336

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Goat Anti Mouse IgG (STAR77...)	HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M365713:200529'

Printed on 19 Aug 2024
