

Datasheet: MCA1778S

BATCH NUMBER 171471

Description:	MOUSE ANTI DOG CD11c
Specificity:	CD11c
Other names:	CD11, INTEGRIN ALPHA X CHAIN
Format:	S/N
Product Type:	Monoclonal Antibody
Clone:	CA11.6A1
Isotype:	IgG1
Quantity:	2 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species

Dog

Species Cross Reactivity

Reacts with: Hooded Seal, Raccoon

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form	Tissue culture supernatant - liquid
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)
RRID	AB_322942
Specificity	<p>Mouse anti Dog CD11c antibody, clone CA11.6A1 recognizes the canine CD11c cell surface antigen, a member of the alpha integrin family. Canine CD11c is expressed by monocytes, granulocytes and by dendritic cells.</p> <p>Mouse anti Dog CD11c, clone CA11.6A1 immunoprecipitates proteins of approximately 95 kDa, corresponding to the common β chain of the CD11/CD18 heterodimer and ~150 kDa, the CD11c; chain from canine leukocyte preparations (Danilenko et al. 1992)</p>
Flow Cytometry	Use 10 μ l of the suggested working dilution to label 10 ⁶ cells or 100 μ l whole blood
References	<ol style="list-style-type: none"> Danilenko, D.M. <i>et al.</i> (1992) Canine leukocyte cell adhesion molecules (LeuCAMs): characterization of the CD11/CD18 family. Tissue Antigens 40: 13-21. McDonough, S.P. and Moore, P.F. (2000) Clinical, hematologic, and immunophenotypic characterization of canine large granular lymphocytosis. Vet Pathol. 37: 637-46. Affolter, V.K. and Moore, P.F. (2002) Localized and disseminated histiocytic sarcoma of dendritic cell origin in dogs. Vet Pathol. 39: 74-83. Catchpole, B. <i>et al.</i> (2002) Generation of blood-derived dendritic cells in dogs with oral malignant melanoma. J Comp Pathol. 126: 238-41. Sanchez, M.A. <i>et al.</i> (2004) Organ-specific immunity in canine visceral leishmaniasis: analysis of symptomatic and asymptomatic dogs naturally infected with <i>Leishmania chagasi</i>. Am J Trop Med Hyg. 70: 618-24. Ibisch, C. <i>et al.</i> (2005) Functional canine dendritic cells can be generated in vitro from peripheral blood mononuclear cells and contain a cytoplasmic ultrastructural marker. J Immunol Methods. 298: 175-82. Mathes, M. <i>et al.</i> (2006) Evaluation of liposomal clodronate in experimental spontaneous autoimmune hemolytic anemia in dogs. Exp Hematol. 34: 1393-402. Isotani, M. <i>et al.</i> (2006) Efficient generation of canine bone marrow-derived dendritic cells. J Vet Med Sci. 68: 809-14. Wang, Y.S. <i>et al.</i> (2007) Characterization of canine monocyte-derived dendritic cells with phenotypic and functional differentiation. Can J Vet Res. 71: 165-74. Liu, C.C. <i>et al.</i> (2008) Transient downregulation of monocyte-derived dendritic-cell differentiation, function, and survival during tumoral progression and regression in an in vivo canine model of transmissible venereal tumor. Cancer Immunol Immunother. 57: 479-91. Bird, R.C. <i>et al.</i> (2008) An allogeneic hybrid-cell fusion vaccine against canine mammary cancer. Vet Immunol Immunopathol. 123: 289-304. Kang, J.W. <i>et al.</i> (2008) Soluble factors-mediated immunomodulatory effects of canine adipose tissue-derived mesenchymal stem cells. Stem Cells Dev. 17: 681-93. Wang, Y.S. <i>et al.</i> (2008) Cytokine profiles of canine monocyte-derived dendritic cells as a function of lipopolysaccharide- or tumor necrosis factor-alpha-induced maturation. Vet Immunol Immunopathol. 118: 186-98.

14. Schwartz, M. *et al.* (2008) Selective CD11a upregulation on neutrophils in the acute phase of steroid-responsive meningitis-arteritis in dogs. [Vet Immunol Immunopathol. 126: 248-55.](#)
15. Ricklin Gutzwiller, M.E. *et al.* (2010) Comparative analysis of canine monocyte- and bone-marrow-derived dendritic cells. [Vet Res. 41: 40.](#)
16. Pai, C.C. *et al.* (2011) Immunopathogenic behaviors of canine transmissible venereal tumor in dogs following an immunotherapy using dendritic/tumor cell hybrid. [Vet Immunol Immunopathol. 139 \(2-4\): 187-99.](#)
17. Figueiredo, M.M. *et al.* (2013) Expression of Toll-like Receptors 2 and 9 in cells of dog jejunum and colon naturally infected with *Leishmania infantum*. [BMC Immunol. 14: 22.](#)
18. Larsen, A.K. *et al.* (2013) Entry and elimination of marine mammal *Brucella* spp. by hooded seal (*Cystophora cristata*) alveolar macrophages *in vitro*. [PLoS One. 8: e70186.](#)
19. Qeska, V. *et al.* (2014) Canine distemper virus infection leads to an inhibitory phenotype of monocyte-derived dendritic cells *in vitro* with reduced expression of co-stimulatory molecules and increased interleukin-10 transcription. [PLoS One. 9 \(4\): e96121.](#)
20. Heinrich, F. *et al.* (2015) Immunophenotyping of immune cell populations in the raccoon (*Procyon lotor*). [Vet Immunol Immunopathol. 168 \(3-4\): 140-6.](#)
21. Paoloni, M. *et al.* (2015) Defining the Pharmacodynamic Profile and Therapeutic Index of NHS-IL12 Immunocytokine in Dogs with Malignant Melanoma. [PLoS One. 10 \(6\): e0129954.](#)
22. Constantinoiu, C.C. *et al.* (2015) Mucosal tolerance of the hookworm *Ancylostoma caninum* in the gut of naturally infected wild dogs. [Parasite Immunol. 37 \(10\): 510-20.](#)
23. Stokol, T. *et al.* (2015) Alkaline phosphatase is a useful cytochemical marker for the diagnosis of acute myelomonocytic and monocytic leukemia in the dog. [Vet Clin Pathol. 44 \(1\): 79-93.](#)
24. Heinrich, F. *et al.* (2015) Passage-dependent morphological and phenotypical changes of a canine histiocytic sarcoma cell line (DH82 cells). [Vet Immunol Immunopathol. 163 \(1-2\): 86-92.](#)
25. Bonnefont-Rebeix, C. *et al.* (2016) Characterization of a novel canine T-cell line established from a spontaneously occurring aggressive T-cell lymphoma with large granular cell morphology. [Immunobiology. 221 \(1\): 12-22.](#)
26. Bird, R.C. *et al.* (2019) Autologous hybrid cell fusion vaccine in a spontaneous intermediate model of breast carcinoma. [J Vet Sci. 20 \(5\): e48.](#)
27. Miguelena Chamorro, B. *et al.* (2023) Characterization of Canine Peyer's Patches by Multidimensional Analysis: Insights from Immunofluorescence, Flow Cytometry, and Single-Cell RNA Sequencing. [Immunohorizons. 7 \(11\): 788-805.](#)
28. Mason, N.J. *et al.* (2021) Development of a fully canine anti-canine CTLA4 monoclonal antibody for comparative translational research in dogs with spontaneous tumors. [MAbs. 13 \(1\): 2004638.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10336 available at: https://www.bio-rad-antibodies.com/SDS/MCA1778S
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M431850:240819'

Printed on 20 May 2026