

Datasheet: MCA1779GA

BATCH NUMBER 161824

Description:	MOUSE ANTI DOG CD11d
Specificity:	CD11d
Other names:	INTEGRIN ALPHA D CHAIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CA11.8H2
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/50 - 1/200
Immunohistology - Frozen (1)	•			
Immunohistology - Paraffin		•		
ELISA			•	
Immunoprecipitation	•			
Western Blotting				

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Dog
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Affinity purified CD11 antigen from canine spleen.
Specificity	Mouse anti Dog CD11d antibody, clone CA11.8H2 recognizes the canine CD11d cell surface antigen, also known as integrin alpha d subunit. The alpha d integrin (CD11d) subunit is expressed as a heterodimer with beta 2 integrin (CD18). CD11d is expressed by a sub-population of macrophages, some CD8 positive T cells and by gamma/delta T cells in the splenic red pulp.
	Over expression of CD11d is noted in histiocytic disorders including hemophagocytic sarcoma, described as a proliferative disorder specifically of CD11+ macrophages (Moore et al. 2006) with macrophages demonstrating marked erythrocytic phagocytosis. Previously, canine histiocytic sarcoma were described as originating entirely from interstitial dendritic cells. Cells from animals with canine hepatosplenic T cell lymphoma (HTSL) strongly express CD11d on malignant cells. Dogs normally demonstrate strong positivity for CD11d on large granular lymphocytes, macrophages and T cells in the splenic red pulp. CD11+ γ / δ T cells constitute ~30% red pulp T cells despite their rarity in peripheral blood, suggesting a splenic red pulp origin of HTSL (Fry et al. 2003).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood
Flow Cytometry References	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood 1. Danilenko, D.M. <i>et al.</i> (1995) A novel canine leukointegrin, alpha d beta 2, is expressed by specific macrophage subpopulations in tissue and a minor CD8+ lymphocyte subpopulation in peripheral blood. J Immunol. 155:35-44. 2. Moore, P.F. <i>et al.</i> (2006) Canine hemophagocytic histiocytic sarcoma: a proliferative disorder of CD11d+ macrophages. Vet Pathol. 43:632-45. 3. Vernau, W. and Moore, P.F. (1999) An immunophenotypic study of canine leukemias and preliminary assessment of clonality by polymerase chain reaction. Vet Immunol Immunopathol. 69:145-64. 4. Fry, M.M. <i>et al.</i> (2003) Hepatosplenic lymphoma in a dog. Vet Pathol. 40:556-62. 5. Stokol, T. <i>et al.</i> (2015) Alkaline phosphatase is a useful cytochemical marker for the diagnosis of acute myelomonocytic and monocytic leukemia in the dog. Vet Clin Pathol. 44 (1): 79-93. 6. Ortiz, A.L. <i>et al.</i> (2015) Gamma delta T-cell large granular lymphocyte lymphoma in a dog. Vet Clin Pathol. 44 (3): 442-7.
	 Danilenko, D.M. et al. (1995) A novel canine leukointegrin, alpha d beta 2, is expressed by specific macrophage subpopulations in tissue and a minor CD8+ lymphocyte subpopulation in peripheral blood. J Immunol. 155:35-44. Moore, P.F. et al. (2006) Canine hemophagocytic histiocytic sarcoma: a proliferative disorder of CD11d+ macrophages. Vet Pathol. 43:632-45. Vernau, W. and Moore, P.F. (1999) An immunophenotypic study of canine leukemias and preliminary assessment of clonality by polymerase chain reaction. Vet Immunol Immunopathol. 69:145-64. Fry, M.M. et al. (2003) Hepatosplenic lymphoma in a dog. Vet Pathol. 40:556-62. Stokol, T. et al. (2015) Alkaline phosphatase is a useful cytochemical marker for the diagnosis of acute myelomonocytic and monocytic leukemia in the dog. Vet Clin Pathol. 44 (1): 79-93. Ortiz, A.L. et al. (2015) Gamma delta T-cell large granular lymphocyte lymphoma in a

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)

Goat Anti Mouse IgG IgA IgM (STAR87...)

RPE

Goat Anti Mouse IgG (STAR76...)

RPE

Rabbit Anti Mouse IgG (STAR13...)

HRP

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) <u>FITC</u>
Goat Anti Mouse IgG (STAR77...) <u>HRP</u>

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M383590:210513'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint