

Datasheet: MCA1783

BATCH NUMBER 159416

Description:	MOUSE ANTI BOVINE INTERFERON GAMMA
Specificity:	IFN GAMMA
Other names:	INTERFERON GAMMA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CC302
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			1/100 - 1/500
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)Membrane permeabilization is required for this application. Bio-Rad recommend the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.

Target Species	Bovine
Species Cross Reactivity	<p>Reacts with: Human, Pig, Dog, Horse, Sheep, Goat, Dolphin, Ferret, Mink, Fin Whale, Rabbit</p> <p>Based on sequence similarity, is expected to react with:Mustelid</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>

Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	<p>UniProt: P07353 Related reagents</p> <p>Entrez Gene: 281237 IFNG Related reagents</p>
RRID	AB_2123454
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
Specificity	<p>Mouse anti Bovine IFNγ antibody, clone CC302 recognizes bovine interferon-gamma, a 143 amino acid cytokine with potent activating, antiviral and anti proliferative properties, produced as a pro-peptide with an additional 23 amino acid N-terminal signal peptide sequence having a molecular weight of ~20 kDa. IFNγ is predominantly secreted by activated T lymphocytes in response to specific mitogens as a result of infection (Rhodes et al. 2000).</p> <p>Mouse anti bovine γ interferon antibody, clone CC302 has been demonstrated to be reactive to a number of mammalian species including human, sheep, dog, pig, goat and mink (Pedersen et al. 2002). Mouse anti Bovine IFNγ antibody, clone CC302 has been used successfully for the evaluation of γ interferon levels in the sera of calves naturally infected with <i>M. avium</i> subsp <i>paratuberculosis</i> (Appana et al. 2013) as a detection reagent using an ELISA.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
ELISA	Biotinylated mouse anti bovine IFNγ, clone CC302 , may be used as the detection reagent in a sandwich ELISA with purified mouse anti bovine IFNγ, clone CC330 , as the capture reagent and recombinant bovine IFNγ as the standard.
References	1. Hasvold, H.J. et al. (2002) <i>In vitro</i> responses to purified protein derivate of caprine T lymphocytes following vaccination with live strains of <i>Mycobacterium avium</i> subsp

- paratuberculosis*. [Vet Immunol Immunopathol. 90 \(1-2\): 79-89.](#)
2. Mwangi, W. *et al.* (2002) DNA-encoded fetal liver tyrosine kinase 3 ligand and granulocyte macrophage-colony-stimulating factor increase dendritic cell recruitment to the inoculation site and enhance antigen-specific CD4⁺ T cell responses induced by DNA vaccination of outbred animals. [J Immunol. 169 \(7\): 3837-46.](#)
 3. Pedersen, L.G. *et al.* (2002) Identification of monoclonal antibodies that cross-react with cytokines from different animal species. [Vet Immunol Immunopathol. 88 \(3-4\): 111-22.](#)
 4. Aasted, B. *et al.* (2002) Cytokine profiles in peripheral blood mononuclear cells and lymph node cells from piglets infected in utero with porcine reproductive and respiratory syndrome virus. [Clin Diagn Lab Immunol. 9 \(6\): 1229-34.](#)
 5. Nielsen, L. *et al.* (2009) Lymphotropism and host responses during acute wild-type canine distemper virus infections in a highly susceptible natural host. [J Gen Virol. 90: 2157-65.](#)
 6. Jaber, J.R. *et al.* (2010) Cross-reactivity of anti-human, anti-porcine and anti-bovine cytokine antibodies with cetacean tissues. [J Comp Pathol. 143: 45-51.](#)
 7. Martel, C.J. & Aasted, B. (2009) Characterization of antibodies against ferret immunoglobulins, cytokines and CD markers. [Vet Immunol Immunopathol. 132:109-15.](#)
 8. Sow, F.B. *et al.* (2011) Respiratory syncytial virus is associated with an inflammatory response in lungs and architectural remodeling of lung-draining lymph nodes of newborn lambs. [Am J Physiol Lung Cell Mol Physiol. 300 \(1\): L12-24.](#)
 9. Ferret-Bernard, S. *et al.* (2011) Mesenteric lymph node cells from neonates present a prominent IL-12 response to CpG oligodeoxynucleotide via an IL-15 feedback loop of amplification. [Vet Res. 42:19.](#)
 10. Lybeck, K.R. *et al.* (2009) Neutralization of interleukin-10 from CD14(+) monocytes enhances gamma interferon production in peripheral blood mononuclear cells from *Mycobacterium avium* subsp. *paratuberculosis*-infected goats. [Clin Vaccine Immunol. 16 \(7\): 1003-11.](#)
 11. Contreras, V. *et al.* (2010) Existence of CD8 α -like dendritic cells with a conserved functional specialization and a common molecular signature in distant mammalian species. [J Immunol. 185: 3313-25.](#)
 12. Fellman, C.L. *et al.* (2011) Cyclosporine A affects the *in vitro* expression of T cell activation-related molecules and cytokines in dogs. [Vet Immunol Immunopathol. 140: 175-80.](#)
 13. Pillet, S. *et al.* (2011) Cellular immune response in the presence of protective antibody levels correlates with protection against 1918 influenza in ferrets. [Vaccine. 29 \(39\): 6793-801.](#)
 14. Jensen, T.H. *et al.* (2009) Early life DNA vaccination with the H gene of Canine distemper virus induces robust protection against distemper. [Vaccine. 27: 5178-83.](#)
 15. Skyberg, J.A. *et al.* (2011) Murine and bovine $\gamma\delta$ T cells enhance innate immunity against *Brucella abortus* infections. [PLoS One. 6:e21978.](#)
 16. Whelan, A.O. *et al.* (2011) Development of an Antibody to Bovine IL-2 Reveals Multifunctional CD4 T(EM) Cells in Cattle Naturally Infected with Bovine Tuberculosis. [PLoS One. 6: e29194.](#)
 17. Costa-Pereira, C. *et al.* (2015) One-year timeline kinetics of cytokine-mediated cellular immunity in dogs vaccinated against visceral leishmaniasis. [BMC Vet Res. 11 \(1\): 92.](#)
 18. Summers, C. *et al.* (2012) The distribution of immune cells in the lungs of classical and atypical ovine pulmonary adenocarcinoma. [Vet Immunol Immunopathol. 146: 1-7.](#)

19. Maślanka T *et al.* (2012) The presence of CD25 on bovine WC1+ $\gamma\delta$ T cells is positively correlated with their production of IL-10 and TGF- β , but not IFN- γ . [Pol J Vet Sci. 15 \(1\): 11-20.](#)
20. Duncombe, L. *et al.* (2013) Investigating the Use of Protein Saver Cards for Storage and Subsequent Detection of Bovine Anti-*Brucella abortus* Smooth Lipopolysaccharide Antibodies and Gamma Interferon. [Clin Vaccine Immunol. 20: 1669-74.](#)
21. Verhelst, D. *et al.* (2014) Parasite distribution and associated immune response during the acute phase of *Toxoplasma gondii* infection in sheep. [BMC Vet Res. 2014 Dec 16;10\(1\):293.](#)
22. Köhler, H. *et al.* (2015) Characterization of a caprine model for the subclinical initial phase of *Mycobacterium avium* subsp. paratuberculosis infection [BMC Veterinary Research. 11 \(1\): 74.](#)
23. Moreira, M.L. *et al.* (2016) Vaccination against canine leishmaniasis increases the phagocytic activity, nitric oxide production and expression of cell activation/migration molecules in neutrophils and monocytes. [Vet Parasitol. 220: 33-45.](#)
24. Rodríguez-Gómez IM *et al.* (2016) Expression of T-bet, Eomesodermin and GATA-3 in porcine $\alpha\beta$ T cells. [Dev Comp Immunol. 60: 115-26.](#)
25. Taylor, G. *et al.* (2015) Efficacy of a virus-vectored vaccine against human and bovine respiratory syncytial virus infections. [Sci Transl Med. 7 \(300\): 300ra127.](#)
26. Moreira, M.L. *et al.* (2015) Cross-reactivity of commercially available anti-human monoclonal antibodies with canine cytokines: establishment of a reliable panel to detect the functional profile of peripheral blood lymphocytes by intracytoplasmic staining. [Acta Vet Scand. 57: 51.](#)
27. El-Naggar, M.M. *et al.* (2015) Development of an improved ESAT-6 and CFP-10 peptide-based cytokine flow cytometric assay for bovine tuberculosis. [Comp Immunol Microbiol Infect Dis. 42: 1-7.](#)
28. McGill, J.L. *et al.* (2016) Vaccination with an Attenuated Mutant of *Ehrlichia chaffeensis* Induces Pathogen-Specific CD4+ T Cell Immunity and Protection from Tick-Transmitted Wild-Type Challenge in the Canine Host. [PLoS One. 11 \(2\): e0148229.](#)
29. Vida, B. *et al.* (2016) Immunologic progression of canine leishmaniasis following vertical transmission in United States dogs. [Vet Immunol Immunopathol. 169: 34-8.](#)
30. Totté, P. *et al.* (2010) CD62L defines a subset of pathogen-specific bovine CD4 with central memory cell characteristics. [Dev Comp Immunol. 34 \(2\): 177-82.](#)
31. Sun, L. *et al.* (2012) The role of proliferation in the regulation of interferon gamma (IFN γ) expression in foals. [Dev Comp Immunol. 36 \(3\): 534-9.](#)
32. Reber, A.J. *et al.* (2006) Evaluation of multiple immune parameters after vaccination with modified live or killed bovine viral diarrhoea virus vaccines. [Comp Immunol Microbiol Infect Dis. 29 \(1\): 61-77.](#)
33. Katepalli, M.P. *et al.* (2008) The effect of age and telomere length on immune function in the horse. [Dev Comp Immunol. 32 \(12\): 1409-15.](#)
34. Hansen, S. *et al.* (2013) Age-related changes in intracellular expression of IFN- γ and TNF- α in equine lymphocytes measured in bronchoalveolar lavage and peripheral blood. [Dev Comp Immunol. 39 \(3\): 228-33.](#)
35. Boshra H *et al.* (2015) A lumpy skin disease virus deficient of an IL-10 gene homologue provides protective immunity against virulent capripoxvirus challenge in sheep and goats. [Antiviral Res. 123: 39-49.](#)
36. Hedges, J.F. *et al.* (2015) Amphotericin B stimulates $\gamma\delta$ T and NK cells, and enhances

- protection from *Salmonella* infection. [Innate Immun. 21 \(6\): 598-608.](#)
37. Johnson, W.C. *et al.* (2008) Bovine WC1(-) gammadeltaT cells incubated with IL-15 express the natural cytotoxicity receptor CD335 (NKp46) and produce IFN-gamma in response to exogenous IL-12 and IL-18. [Dev Comp Immunol. 32 \(8\): 1002-10.](#)
38. Dewals, B.G., *et al.* (2011) Malignant catarrhal fever induced by Alcelaphine herpesvirus 1 is characterized by an expansion of activated CD3+CD8+CD4- T cells expressing a cytotoxic phenotype in both lymphoid and non-lymphoid tissues [Vet Res. 42\(1\): 95.](#)
39. Maggioli, M.F. *et al.* (2016) Increased TNF- α /IFN- γ /IL-2 and Decreased TNF- α /IFN- γ Production by Central Memory T Cells Are Associated with Protective Responses against Bovine Tuberculosis Following BCG Vaccination. [Front Immunol. 7: 421.](#)
40. Cassady-cain, R.L. *et al.* (2017) Inhibition of Antigen-Specific and Nonspecific Stimulation of Bovine T and B Cells by Lymphostatin from Attaching and Effacing *Escherichia coli*. [Infect Immun. 85 \(2\)Jan 26 \[Epub ahead of print\].](#)
41. Wattedegera, S.R. *et al.* (2017) Enhancing the toolbox to study IL-17A in cattle and sheep. [Vet Res. 48 \(1\): 20.](#)
42. DaSilva, A.V.A. *et al.* (2018) Morphophysiological changes in the splenic extracellular matrix of *Leishmania infantum*-naturally infected dogs is associated with alterations in lymphoid niches and the CD4+ T cell frequency in spleens. [PLoS Negl Trop Dis. 12 \(4\): e0006445.](#)
43. Higgins, J.L. *et al.* (2018) Cell mediated immune response in goats after experimental challenge with the virulent *Brucella melitensis* strain 16M and the reduced virulence strain Rev. 1. [Vet Immunol Immunopathol. 202: 74-84.](#)
44. Roos, E.O. *et al.* (2018) IP-10: A potential biomarker for detection of Mycobacterium bovis infection in warthogs (*Phacochoerus africanus*). [Vet Immunol Immunopathol. 201: 43-8.](#)
45. Aguiar-Soares, R.D.O. *et al.* (2020) Phase I and II Clinical Trial Comparing the LBSap, Leishmune[®], and Leish-Tec[®] Vaccines against Canine Visceral Leishmaniasis. [Vaccines \(Basel\). 8 \(4\)Nov 17 \[Epub ahead of print\].](#)
46. Fedorka, C.E. *et al.* (2019) Alteration of the mare's immune system by the synthetic progestin, altrenogest. [Am J Reprod Immunol. 82 \(2\): e13145.](#)
47. Lacasta, A. *et al.* (2021) Synergistic Effect of Two Nanotechnologies Enhances the Protective Capacity of the *Theileria parva* Sporozoite p67C Antigen in Cattle. [J Immunol. Jan 08 \[Epub ahead of print\].](#)
48. Arrieta-Villegas, C. *et al.* (2020) Immunogenicity and Protection against *Mycobacterium caprae* Challenge in Goats Vaccinated with BCG and Revaccinated after One Year. [Vaccines \(Basel\). 8 \(4\): 751.](#)
49. Witonsky, S. *et al.* (2019) Can levamisole upregulate the equine cell-mediated macrophage (M1) dendritic cell (DC1) T-helper 1 (CD4 Th1) T-cytotoxic (CD8) immune response *in vitro*? [J Vet Intern Med. 33 \(2\): 889-96.](#)
50. Freer, H. *et al.* (2017) A monoclonal antibody for detection of intracellular and secreted interleukin-2 in horses. [Vet Immunol Immunopathol. 191: 30-5.](#)
51. Rodrigues, V. *et al.* (2017) Development of a bead-based multiplexed assay for simultaneous quantification of five bovine cytokines by flow cytometry. [Cytometry A. 91 \(9\): 901-7.](#)
52. Hillmann, A. *et al.* (2019) A novel direct co-culture assay analyzed by multicolor flow cytometry reveals context- and cell type-specific immunomodulatory effects of equine

mesenchymal stromal cells. [PLoS One. 14 \(6\): e0218949.](#)

53. Villa-Mancera, A. *et al.* (2021) Phage display-based vaccine with cathepsin L and excretory-secretory products mimotopes of *Fasciola hepatica*. induces protective cellular and humoral immune responses in sheep. [Vet Parasitol. 289: 109340.](#)

54. Risalde, M.Á. *et al.* (2017) Development and evaluation of an interferon gamma assay for the diagnosis of tuberculosis in red deer experimentally infected with *Mycobacterium bovis*. [BMC Vet Res. 13 \(1\): 341.](#)

55. Bidart, J. *et al.* (2020) A New Cage-Like Particle Adjuvant Enhances Protection of Foot-and-Mouth Disease Vaccine. [Front Vet Sci. 7: 396.](#)

56. Damani-Yokota, P. *et al.* (2021) Transcriptional programming and gene regulation in WC1⁺ γδ T cell subpopulations. [Mol Immunol. 142: 50-62.](#)

57. Ciliberti, M.G. *et al.* (2022) Green extraction of bioactive compounds from wine lees and their bio-responses on immune modulation using *in vitro*. sheep model. [J Dairy Sci. Mar 17 \[Epub ahead of print\].](#)

Further Reading 1. Rhodes, S. *et al.* (2000) Distinct response kinetics of gamma interferon and interleukin-4 in bovine tuberculosis. [Infect Immun. 68:5393-400.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1783>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M383156:210513'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)