

Datasheet: MCA1784

BATCH NUMBER 147834

Description:	RAT ANTI HUMAN EGF RECEPTOR
Specificity:	EGF R
Other names:	EPIDERMAL GROWTH FACTOR RECEPTOR
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	ICR10
Isotype:	IgG2a
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting		▪		
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Extracellular domain of human EGF-receptor from head and neck carcinoma.
External Database Links	<p>UniProt: P00533 Related reagents</p> <p>Entrez Gene: 1956 EGFR Related reagents</p>
Synonyms	ERBB1
RRID	AB_322787
Specificity	<p>Rat anti Human EGF Receptor antibody, clone ICR10 recognizes the human epidermal growth factor receptor (EGF-R), which is over expressed in a high proportion of breast cancer cells and in a range of other carcinomas.</p> <p>High level expression of EGFR is often associated with advanced disease and poor prognosis.</p> <p>Rat anti Human EGF Receptor antibody, clone ICR10 binds to epitope B from EGFR (Lottaz et al. 2010 and Modjtahedi et al. 1993) and has an affinity of 6.7×10^{-9} M.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10^6 cells in 100ul.
Histology Positive Control Tissue	Breast Carcinoma
References	<ol style="list-style-type: none"> Lottaz,C. <i>et al.</i> (2010) Transcriptional Profiles of CD133+ and CD133- Glioblastoma-Derived Cancer Stem Cell Lines Suggest Different Cells of Origin Cancer Res. 70: 2030-40. Modjtahedi, H. <i>et al.</i> (1993) Antitumor activity of combinations of antibodies directed against different epitopes on the extracellular domain of the human EGF receptor. Cell Biophys. 22 (1-3): 129-46. Modjtahedi, H. <i>et al.</i> (2012) Immunohistochemical discrimination of wild-type EGFR from EGFRvIII in fixed tumour specimens using anti-EGFR mAbs ICR9 and ICR10. Br J Cancer. 106 (5): 883-8. Grinberg, O. <i>et al.</i> (2013) Antibody modified Bovine Serum Albumin microspheres for targeted delivery of anticancer agent Gemcitabine Polymers for Advanced Technologies. 24 (3): 294-299. Gilcrease, M.Z. <i>et al.</i> (2009) Alpha6beta4 integrin crosslinking induces EGFR clustering and promotes EGF-mediated Rho activation in breast cancer. J Exp Clin Cancer Res. 28: 67. Tilburgs, T. <i>et al.</i> (2015) Human HLA-G+ extravillous trophoblasts: Immune-activating

cells that interact with decidual leukocytes. [Proc Natl Acad Sci U S A. 112 \(23\): 7219-24.](#)
7. Tilburgs, T. *et al.* (2015) The HLA-G cycle provides for both NK tolerance and immunity at the maternal-fetal interface. [Proc Natl Acad Sci U S A. 112 \(43\): 13312-7.](#)
8. Khelwatty, S.A. *et al.* (2015) Acquired resistance to anti-EGFR mAb ICR62 in cancer cells is accompanied by an increased EGFR expression, HER-2/HER-3 signalling and sensitivity to pan HER blockers. [Br J Cancer. 113 \(7\): 1010-9.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1784>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M373432:200911'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)