

Datasheet: MCA1784F

BATCH NUMBER 154597

Description:	RAT ANTI HUMAN EGF RECEPTOR:FITC
Specificity:	EGF R
Other names:	EPIDERMAL GROWTH FACTOR RECEPTOR
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	ICR10
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative	0.09% Sodium Azide						
Stabilisers	1% Bovine Serum Albumin						
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml						

Immunogen	Extracellular domain of human EGF-receptor from head and neck carcinoma.
External Database Links	<p>UniProt: P00533 Related reagents</p> <p>Entrez Gene: 1956 EGFR Related reagents</p>
Synonyms	ERBB1
RRID	AB_324284
Specificity	<p>Rat anti Human EGF Receptor antibody, clone ICR10 recognizes the human epidermal growth factor receptor (EGF-R), which is over expressed in a high proportion of breast cancer cells and in a range of other carcinomas.</p> <p>High level expression of EGFR is often associated with advanced disease and poor prognosis.</p> <p>Rat anti Human EGF Receptor antibody, clone ICR10 binds to epitope B from EGFR (Lottaz et al. 2010 and Modjtahedi et al. 1993) and has an affinity of 6.7×10^{-9} M.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10^6 cells in 100ul.
References	<ol style="list-style-type: none"> Lottaz, C. <i>et al.</i> (2010) Transcriptional Profiles of CD133+ and CD133- Glioblastoma-Derived Cancer Stem Cell Lines Suggest Different Cells of Origin Cancer Res. 70: 2030-40. Modjtahedi, H. <i>et al.</i> (1993) Antitumor activity of combinations of antibodies directed against different epitopes on the extracellular domain of the human EGF receptor. Cell Biophys. 22 (1-3): 129-46. Modjtahedi, H. <i>et al.</i> (2012) Immunohistochemical discrimination of wild-type EGFR from EGFRvIII in fixed tumour specimens using anti-EGFR mAbs ICR9 and ICR10. Br J Cancer. 106 (5): 883-8. Grinberg, O. <i>et al.</i> (2013) Antibody modified Bovine Serum Albumin microspheres for targeted delivery of anticancer agent Gemcitabine Polymers for Advanced Technologies. 24 (3): 294-299. Gilcrease, M.Z. <i>et al.</i> (2009) Alpha6beta4 integrin crosslinking induces EGFR clustering and promotes EGF-mediated Rho activation in breast cancer. J Exp Clin Cancer Res. 28: 67. Tilburgs, T. <i>et al.</i> (2015) Human HLA-G+ extravillous trophoblasts: Immune-activating cells that interact with decidual leukocytes. Proc Natl Acad Sci U S A. 112 (23): 7219-24. Tilburgs, T. <i>et al.</i> (2015) The HLA-G cycle provides for both NK tolerance and immunity at the maternal-fetal interface. Proc Natl Acad Sci U S A. 112 (43): 13312-7. Khelwatty, S.A. <i>et al.</i> (2015) Acquired resistance to anti-EGFR mAb ICR62 in cancer cells is accompanied by an increased EGFR expression, HER-2/HER-3 signalling and sensitivity to pan HER blockers. Br J Cancer. 113 (7): 1010-9.

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1784F>
10041

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:FITC \(MCA6005F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M395619:220519'

Printed on 22 Mar 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)