

Datasheet: MCA1799 BATCH NUMBER 155425

Description:	MOUSE ANTI HUMAN ESTROGEN RECEPTOR ALPHA		
Specificity:	ESTROGEN RECEPTOR ALPHA		
Format:	S/N		
Product Type:	Monoclonal Antibody		
Clone:	6F11		
Isotype:	lgG1		
Quantity:	1 ml		

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>						
	rad-antibodies.com/proto	rad-antibodies.com/protocols.					
	Flow Cytometry	Yes	No	Not Determined	Suggested Dilution		
	Immunohistology - Frozen (1)	•			1/40 - 1/60		
	Immunohistology - Paraffin (2)	•			1/40 - 1/80		
	ELISA						
	Immunoprecipitation						
	Western Blotting				1/50 - 1/100		
	necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls. (1)The use of Zamboni's fixative is recommended for best results (2)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.Sodium citrate buffer pH 6.0 is recommended for this purpose.						
Target Species	Human						
Product Form	Tissue Culture Supernatant - liquid						
Preservative Stabilisers	0.09% Sodium Azide						
Immunogen	Recombinant human estrogen receptor (alpha form).						

External Database Links	UniProt: <u>P03372</u> <u>Related reagents</u>			
	Entrez Gene:			
	2099 ESR1 <u>Related reagents</u>			
Synonyms	ESR, NR3A1			
RRID	AB_2102070			
Fusion Partners	Spleen cells from immunised mice were fused with cells of the mouse p3-NS1-Ag4-1 myeloma cell line.			
Specificity	Mouse anti Human estrogen alpha antibody, clone 6F11 recognizes the human estrogen receptor alpha chain (ER α), also known as the estradiol receptor or nuclear receptor subfamily 3 group A member 1. ER α is a ~65 kDa steroid hormone receptor containing an N-terminal (AF-1) ligand independent transactivation domain, a DNA binding domain and a C-terminal ligand binding domain which overlaps with an (AF-2) domain. ER α binds to DNA as a homodimer (Klinge 2001) and can also form heterodimers with Estrogen receptor beta.			
	The detection of estrogen (ER) and progesterone (PR) receptors using immunohistochemical staining of formalin fixed, paraffin embedded (FFPE) tissue, has gradually replaced ligand binding assays, to become the most common method for the determination of the ER/PR status of breast tumors (<u>Yaziji <i>et al.</i> 2008</u>). Approximately 75% to 80% of breast tumors have estrogen and/or progesterone receptors, and the presence of these receptors helps determine both the patients prognosis and the effectiveness of hormonal therapy (<u>Bhargava <i>et al.</i> 2012</u>).			
	Mouse anti human estrogen alpha antibody, clone 6F11 has been used successfully for identification of ER α on breast cancer cell lysates using Western blotting and for the immunohistochemical detection of ER α in breast cancer tissues (<u>Ambroise <i>et al.</i> 2011</u>).			
Histology Positive Control Tissue	Breast carcinoma			
References	 Bevitt, D.J. <i>et al.</i> (1997) New monoclonal antibodies to oestrogen and progesterone receptors effective for paraffin section immunohistochemistry. <u>J Pathol. 183 (2): 228-32.</u> Jongen, V. <i>et al.</i> (2009) Expression of estrogen receptor-alpha and -beta and progesterone receptor-A and -B in a large cohort of patients with endometrioid endometrial cancer. <u>Gynecol Oncol. 112: 537-42.</u> Ambroise, M. <i>et al.</i> (2011) Immunohistochemical profile of breast cancer patients at a tertiary care hospital in South India. <u>Asian Pac J Cancer Prev. 12: 625-9.</u> Droog, M. <i>et al.</i> (2017) Estrogen receptor α wields treatment-specific enhancers between morphologically similar endometrial tumors. <u>Proc Natl Acad Sci U S A. 114 (8): E1316-E1325.</u> Krishnaswamy, U. <i>et al.</i> (2013) Correlation of Her-2 neu over-expression with clinico 			

	pathological features of carcinoma breast Apollo Medicine. 10 (4): 313-7.
Storage	Store at +4°C or at -20°C if preferred.
	Storage in frost-free freezers is not recommended.
	This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend
	microcentrifugation before use.
Guarantee	Guaranteed until date of expiry. Please see product label.
Health And Safety	Material Safety Datasheet documentation #10053 available at:
Information	https://www.bio-rad-antibodies.com/SDS/MCA1799
	10053
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12)	RPE			
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>				
Goat Anti Mouse IgG (STAR76)	RPE			
Rabbit Anti Mouse IgG (STAR13)	HRP			
Goat Anti Mouse IgG (STAR70)	FITC			
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,			
	DyLight®650, DyLight®680, DyLight®800,			
	<u>FITC</u> , <u>HRP</u>			
Rabbit Anti Mouse IgG (STAR9)	FITC			
Goat Anti Mouse IgG (STAR77)	HRP			
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP			

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	d.com	Email: antibody_sales_uk@bio-ra	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M361300:200210'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint